

# Researchers create tool to predict kidney failure or death after injury

September 2 2013

---

Researchers from Brigham and Women's Hospital (BWH) have developed a risk score calculation that can help predict which patients with rhabdomyolysis (a condition that occurs due to muscle damage) may be at risk for the severe complication of kidney failure or death. This research will publish online September 2, 2013 in *JAMA Internal Medicine*.

Rhabdomyolysis occurs when muscles are crushed and rupture, leaking [toxic compounds](#) into the circulation and can be caused by any condition that damages skeletal muscle and causes injury. Risk factors include crush injuries, alcoholism, [heat stroke](#) or severe exertion including [strenuous exercise](#) and as a rare side effect of taking cholesterol lowering medications called statins. One of the compounds called myoglobin can get caught in the kidneys and cause kidney failure and even death in severe cases.

"Currently doctors cannot easily predict which patients are most likely to have severe [kidney failure](#) or die from rhabdomyolysis," explained Gearoid McMahon, MB, BCh, a clinical fellow in BWH's Department of Medicine and lead author on this study. "Using routinely available clinical variables, we have developed a new risk prediction score that doctors can use to help predict a patient's expected outcome and plan for treatment accordingly."

The researchers conducted a [retrospective cohort study](#) of 2,371 patients admitted between 2000 and 2011 and analyzed variables that are thought

to be associated with poor outcomes in this patient population, such as age, gender, and CPK levels. Using data from this analysis, researchers created a [risk prediction](#) score based on the variables that were most significantly associated with poor outcomes. The final variables that were included in the created model were age, gender, levels of initial phosphate, calcium, creatinine and CO2, CPK, and cause of rhabdomyolysis.

"This model may be particularly useful to evaluate and triage patients in the Emergency Department (ED)," said Sushrut Waikar, MD, MPH, director of Renal and Ambulatory Services at BWH and senior author of the study. "By current practice standards, patients with rhabdomyolysis are treated similarly regardless of the underlying cause and overall risk profile. Knowledge of the predicted risk of adverse outcomes may lead clinicians to choose different treatment options such as intravenous fluid administration in the ED followed by discharge with plans for repeat outpatient labs, rather than inpatient hospitalization for observation."

The researchers note that the next step will be to validate this risk score in other populations and to test its ability to guide treatment decisions. An online version of the Brigham rhabdomyolysis [risk score](#) is available at: <http://www.brighamandwomens.org/research/rhabdo/default.aspx>.

This research is consistent with the recent trend in medicine toward patient-centered outcomes research (PCOR), which focuses on identifying the risk of each patient and tailoring the treatment to improve their individual outcomes. BWH recently established the Patient-centered Comparative Effectiveness Research Center (PCERC) for researchers, like McMahon, that are dedicated to improving the health outcomes of BWH patients.

Provided by Brigham and Women's Hospital

Citation: Researchers create tool to predict kidney failure or death after injury (2013, September 2) retrieved 10 April 2024 from <https://medicalxpress.com/news/2013-09-tool-kidney-failure-death-injury.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.