


USPSTF: evidence lacking for ankle brachial index screening

September 3 2013



The U.S. Preventive Services Task Force has found that there is currently insufficient evidence to weigh the benefits and harms of use of the ankle-brachial index for screening for peripheral artery disease and for predicting cardiovascular disease risk, according to an evidence review and recommendation statement published in the Sept. 3 issue of the *Annals of Internal Medicine*.

(HealthDay)—The U.S. Preventive Services Task Force (USPSTF) has found that there is currently insufficient evidence to weigh the benefits and harms of use of the ankle-brachial index (ABI) for screening for peripheral artery disease and for predicting cardiovascular disease risk, according to an evidence review and recommendation statement published in the Sept. 3 issue of the *Annals of Internal Medicine*.

Jennifer S. Lin, M.D., and colleagues from Kaiser Permanente Northwest in Portland, Ore., reviewed the evidence on the ability of ABI to predict cardiovascular disease morbidity and mortality, independent

of the Framingham Risk Score (FRS). Based on one large meta-analysis involving 43,919 participants, the researchers found that ABI could reclassify 10-year [coronary artery disease](#) risk, but did not report measures of appropriate reclassification (net reclassification improvement [NRI]). In four heterogeneous risk prediction studies, the magnitude of the NRI was likely small for predicting coronary artery disease or cardiovascular disease events when ABI was added to FRS.

Based of these findings, Virginia A. Moyer, M.D., M.P.H., and colleagues from the USPSTF, updated the 2005 recommendation on screening for peripheral artery disease. The researchers concluded that the current evidence is insufficient to assess the balance of benefits and harms associated with ABI screening for [peripheral artery disease](#) and cardiovascular risk assessment in adults (I Statement).

"Even though it is not clear if the ABI helps health care professionals make treatment decisions, there are many things clinicians can do to help their patients reduce their chance of heart attacks and strokes," USPSTF co-vice chair Albert Siu, M.D., M.S.P.H., said in a statement. "The USPSTF recommends screening for and managing high blood pressure, [high blood cholesterol](#), and diabetes and screening for tobacco use and providing cessation support."

More information: [Full Text - Evidence Review](#)
[Full Text - Recommendation Statement](#)
[Editorial](#)

Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: USPSTF: evidence lacking for ankle brachial index screening (2013, September 3)
retrieved 2 May 2024 from
<https://medicalxpress.com/news/2013-09-uspstf-evidence-lacking-ankle-brachial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.