

Women in lower green space areas show higher overall levels of stress, new research shows

September 17 2013



The new study shows that there are significant gender differences in stress patterns by levels of green space.

(Medical Xpress)—Women living in deprived areas with little green space are more likely to be stressed than men living in the same circumstances, according to research published this week in an international journal on public health.

Gender differences in stress patterns



While contact with green space in deprived areas is associated with benefits to mental health, this new study shows that there are significant gender differences in stress patterns by levels of green space.

Women in lower green space areas show higher overall <u>levels of stress</u>, according to the research, led by OPENspace research centre at the Universities of Edinburgh and Heriot-Watt, working in collaboration with the Universities of Glasgow and Westminster, Biomathematics and Statistics Scotland and the James Hutton Institute. The same does not appear to be true of men living in the same areas; an anomaly which the study suggests requires further investigation.

Researchers looked specifically at the concentration of cortisol, the stress hormone, in men and women living in deprived <u>urban areas</u> in Scotland as well as people's perception of their stress levels.

They measured the relationship between gender and percentage of green space on mean cortisol concentrations and found that there was a positive effect of higher green space on women, but not in men.

The effects of contact with green space and a lowering of stress levels is thought to be associated with factors including increased <u>physical</u> <u>activity</u> which improves mood; increased social contact and better mental wellbeing. Contact with nature has also been shown to have positive effects on <u>blood pressure</u> and <u>heart rate</u>.

However, most studies which have measured cortisol levels in relation to contact with nature have focused only on the levels immediately before and after contact with nature.

New study



This new study measured the daily (diurnal) patterns in relation to the long-term effects of familiar, everyday environments, set within the context of people's normal patterns of activity and experience. The study concludes:

- in both men and women, perceived stress was higher in low green space areas, but women's perceived stress was significantly higher in low green space areas than men's
- perceived stress was higher for people with no garden, especially men
- both men and women living in deprived areas with higher levels of green space report less perceived stress and appear to be more resilient to the negative effects of urban deprivation

Speaking on behalf of the research team, lead author Dr Jenny Roe, Heriot Watt University said, "These results are important in understanding how neighbourhood green space might contribute to public health improvement. Stress is known to impact on cardiovascular health, alongside other risk factors such as genetics, age, diet and physical activity, but little is known about the contributions of environmental factors.

"We already know that higher levels of green space are associated with reduced cardio-vascular mortality. Our new study indicates that neighbourhood green space is associated with perceptions of stress as well as the levels of <u>stress hormones</u> in the body and this may be a pathway by which the environment can impact health.

"While we need more research to understand these mechanisms, our study represents a valuable step in establishing a biological pathway linking green space with <u>stress levels</u> in deprived urban environments."



Provided by Heriot-Watt University

Citation: Women in lower green space areas show higher overall levels of stress, new research shows (2013, September 17) retrieved 30 April 2024 from https://medicalxpress.com/news/2013-09-women-green-space-areas-higher.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.