

Atherosclerosis in HIV patients linked to infection, not treatment

October 30 2013

HIV infection, not antiretroviral therapy (ART), is associated with risk for atherosclerosis in patients with no history of smoking, particularly those infected for eight years or more. Results of the study led by Moïse Desvarieux, MD, PhD, at Columbia University's Mailman School of Public Health and at the Institut National de la Santé et de la Recherche Médicale (INSERM), are published in the latest issue of the journal *AIDS*.

A growing body of literature reports that HIV patients have higher risk for atherosclerotic disease and cardiovascular disease, but it has been unknown whether HIV infection or ART are the responsible factors, and to what extent traditional cardiovascular risk factors, chiefly smoking, are responsible. Previous studies on HIV, ART, and atherosclerosis had included smokers which made it difficult to disentangle the relationship. While previous studies had looked at specific inflammatory markers, none had studied the impact of inflammatory imbalance on atherosclerosis in the context of HIV. The authors hypothesized that the imbalance between pro- and anti-inflammatory markers would matter more for atherosclerosis than specific markers.

To explore the question, the researchers performed a retrospective study of 100 HIV-positive never-smokers for a minimum of 2 years, of whom half had never received ART and half had been treated with ART for at least 4 years, along with 50 HIV-negative controls. They looked for correlations between measures of carotid intima-media thickness, or c-IMT—a measure of cardiovascular health—and, in a first, inflammatory

imbalance. Statistical adjustments were made for age, hypertension, and diabetes as well as HIV-infected patients with extremely low counts of lymphocytes, a type of white blood cell.

They found levels of c-IMT were significantly elevated in patients who had been infected for eight years or longer, even those not on ART. Longer time with the infection was also associated with weakened anti-inflammatory response profiles, irrespective of the pro-inflammatory response. "Lower anti-inflammatory response has been linked to increased risk for atherosclerosis, pointing to a potential mechanism within the context of HIV," notes Dr. Franck Boccard, professor of cardiology at Saint-Antoine hospital in Paris and a co-author of the study who will be initiating a 4-month sabbatical at Mailman this week.

The study does not determine whether HIV infection is a cause of atherosclerosis, the authors caution.

But it does contribute significantly by the careful choice of never-smokers and the hard-to-attain group of subjects with 2-year long HIV infection who had not yet required ART initiation, thus allowing for the study of HIV-[infection](#) alone. "We suggest that larger studies incorporate and validate the importance of inflammatory imbalance in occurrence of [atherosclerosis](#)" concludes Dr. Desvarieux, Mailman School associate professor of Epidemiology, who led the transnational multidisciplinary collaboration with scientists at INSERM and the Paris hospital systems.

About 1.1 million Americans live with HIV, according to the Centers for Disease Control. Globally, 34 million are infected, according to the World Health Organization.

Provided by Columbia University's Mailman School of Public Health

Citation: Atherosclerosis in HIV patients linked to infection, not treatment (2013, October 30)
retrieved 27 June 2024 from <https://medicalxpress.com/news/2013-10-atherosclerosis-hiv-patients-linked-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.