


# Bowel cancer patients miss out on life-saving liver surgery

October 29 2013

---



Bowel cancer patients whose disease has spread to the liver are not being referred to specialists for potentially life-saving surgery, according to research published in the *British Journal of Surgery*.

More than 40,000 people are diagnosed with [bowel cancer](#) every year and the disease spreads to the liver in around half of all cases. But this Cancer Research UK led study, which looked at rates of liver [surgery](#) in Yorkshire, found that only a fifth of patients were being referred to liver specialists.

Referral rates also varied widely between the seven hospitals in the area. While one hospital referred as many as 43 per cent of patients on to see liver specialists, another referred as few as 13 per cent.

Researchers at St James's University Hospital, Leeds, looked at the treatment of more than 600 patients with bowel cancer that had spread to the liver. Around one in five were referred to liver specialists and, of these patients, three quarters were deemed to be fit for liver surgery.

Patients who the bowel specialists decided shouldn't undergo a liver operation, despite being fit for surgery, were reassessed by the liver specialists using the same information. In 44 per cent of cases the liver specialists disagreed.

All patients are seen by bowel cancer specialists, who can refer them to liver specialists if they think they might benefit from liver treatment. The most common treatment is surgery to remove any tumours from the liver – which may help cure the patient.

Mr Alastair Young, study author and surgical registrar based at St James's University Hospital, said: "Bowel cancer specialists are deciding whether or not a patient is suitable for liver surgery. And we're seeing liver specialists disagree with the decision in almost half of cases, meaning that some patients are denied an operation that could save their lives.

"One of the striking things about this work is that our region has been shown to have the highest rates in the country of liver surgery for bowel [cancer patients](#). But we've shown a significant gap between current and best practice in our region, suggesting there is plenty of room for improvement up and down the country.

"These decisions are not made lightly but [liver](#) and bowel specialists need to talk to each other much more to ensure every patient has the best possible chance of beating the disease and to reduce the worrying difference in referral rates."

Professor Peter Johnson, Cancer Research UK's chief clinician, said:  
"This shows how important it is for cancer specialists to collaborate when discussing treatment so all patients can benefit from their collective medical expertise.

"The patient's welfare is key so the benefits of surgery must be weighed against the risks. Not all [patients](#) will be eligible for these operations because their cancer may be too advanced, they may have other health problems or they may not want to have surgery."

**More information:** Young, A. L., et al, Variation in referral practice for patients with colorectal cancer liver metastases, *British Journal of Surgery*, 2013. [DOI: 10.1002/bjs.9285](https://doi.org/10.1002/bjs.9285)

Provided by Cancer Research UK

Citation: Bowel cancer patients miss out on life-saving liver surgery (2013, October 29) retrieved 16 August 2024 from <https://medicalxpress.com/news/2013-10-bowel-cancer-patients-life-saving-liver.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--