

Brain development differs in children who stutter

October 10 2013

(Edmonton) A new study by a University of Alberta researcher shows that children who stutter have less grey matter in key regions of the brain responsible for speech production than children who do not stutter.

The findings not only improve our understanding of how the brain is built for [speech](#) production and why people stutter, but also affirm the importance of seeking treatment early, using approaches such as those pioneered by the Institute for Stuttering Treatment and Research in the Faculty of Rehabilitation Medicine at the U of A, said Deryk Beal, ISTAR's executive director.

Previous research has used MRI scans to look at structural differences between the brains of adults who stutter and those who do not. The problem with that approach is the scans come years after the onset of stuttering, typically between the ages of two and five years, Beal said.

"You can never be quite sure whether the differences in brain structure or function you're looking at were the result of a lifetime of coping with a speech disorder or whether those [brain differences](#) were there from the beginning," explained Beal, a speech-language pathologist.

For his study, Beal scanned the brains of 28 children ranging from five to 12 years old. Half the children were diagnosed with stuttering; the other half served as a control.

Results showed that the inferior frontal gyrus region of the brain

develops abnormally in children who stutter. This is important because that part of the brain is thought to control articulatory coding—taking information our [brain](#) understands about language and sounds and coding it into speech movements.

"If you think about the characteristics of stuttering—repetitions of the first sounds or syllables in a word, prolongation of sounds in a word—it's easy to hypothesize that it's a speech-motor-control problem," explained Beal. "The type of stuttering treatment we deliver at ISTAR is delivered with this limitation of the speech system in mind, and we have good success in stuttering treatment."

Beal initiated the research at the University of Toronto and completed the work upon his arrival at the U of A. He sees the results as a first step toward testing to see how [grey matter](#) volumes are influenced by stuttering treatment and understanding motor-sequence learning differences between [children](#) who [stutter](#) and those who do not.

"The more we know about motor learning in these kids, the more we can adjust our treatment—deliver it in a shorter period of time, deliver it more effectively."

More information: The study was published in the September issue of the peer-reviewed journal *Cortex*.

Provided by University of Alberta

Citation: Brain development differs in children who stutter (2013, October 10) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-10-brain-differs-children-stutter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.