

Coexisting pain tied to worse knee replacement outcomes

October 26 2013



(HealthDay)—Musculoskeletal pain in the low back, hips, and the nonoperatively treated knee before total knee replacement is associated with poorer physical function at six months after surgery, according to a study published in the Oct. 16 issue of *The Journal of Bone & Joint Surgery*.

David C. Ayers, M.D., from the University of Massachusetts Medical School in Worcester, and colleagues examined the correlation of bilateral knee and hip pain and low back pain before primary unilateral total knee replacement with physical function at six months after surgery using data from a prospective cohort of 180 patients (61 percent women; mean age, 65.1 years).

The researchers found that patients with a greater number of



preoperative locations of moderate to severe pain had a lower mean Short Form-36 (SF-36) physical component summary score at six months. Moderate to severe preoperative pain in the contralateral knee, ipsilateral and contralateral hip, and low back were all significantly associated with poorer function at six months after total knee replacement, after adjustment for age, sex, body mass index, and SF-36 mental component summary score.

"These data suggest that clinicians, policy makers, and researchers may choose to quantify the total burden of musculoskeletal pain in the knees, <a href="https://hips.nih.gov/hips.nih.go

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2013 HealthDay. All rights reserved.

Citation: Coexisting pain tied to worse knee replacement outcomes (2013, October 26) retrieved 4 May 2024 from

https://medicalxpress.com/news/2013-10-coexisting-pain-tied-worse-knee.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.