

# Large study to examine if vitamin D prevents diabetes

October 22 2013

---

Researchers have begun the first definitive, large-scale clinical trial to investigate if a vitamin D supplement helps prevent or delay type 2 diabetes in adults who have prediabetes, who are at high risk for developing type 2. Funded by the National Institutes of Health, the study is taking place at about 20 study sites across the United States.

The multiyear Vitamin D and Type 2 Diabetes (D2d) study will include about 2,500 people. Its goal is to learn if vitamin D—specifically D3 (cholecalciferol)—will prevent or delay type 2 [diabetes](#) in adults aged 30 or older with [prediabetes](#). People with prediabetes have blood glucose levels that are higher than normal but not high enough to be called diabetes.

"This study aims to definitively answer the question: Can vitamin D reduce the risk of developing [type 2 diabetes](#)?" said Myrlene Staten, M.D., D2d project officer at the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), part of NIH. "Vitamin D use has risen sharply in the U.S. in the last 15 years, since it has been suggested as a remedy for a variety of conditions, including prevention of type 2 diabetes. But we need rigorous testing to determine if vitamin D will help prevent diabetes. That's what D2d will do."

"Past observational studies have suggested that higher levels of vitamin D may be beneficial in preventing type 2 diabetes, but until this large, randomized and controlled clinical trial is complete, we won't know if taking vitamin D supplements lowers the risk of diabetes," said

Anastassios G. Pittas, M.D., the study's principal investigator at Tufts Medical Center, Boston.

D2d is the first study to directly examine if a daily dose of 4,000 International Units (IUs) of vitamin D—greater than a typical adult intake of 600-800 IUs a day, but within limits deemed appropriate for clinical research by the Institute of Medicine—helps keep people with prediabetes from getting type 2 diabetes. Based on observations from earlier studies, researchers speculate that vitamin D could reduce the diabetes risk by 25 percent. The study will also examine if sex, age or race affect the potential of vitamin D to reduce [diabetes risk](#).

"An estimated 79 million Americans have prediabetes, and nearly 26 million more have diabetes," said NIDDK Director Griffin P. Rodgers, M.D. "With D2d, we seek evidence for an affordable and accessible way to help prevent or delay type 2 diabetes."

Researchers are recruiting volunteers to take part in D2d. Half of the participants will receive vitamin D. The other half will receive a placebo—a pill that has no drug effect. Participants will have check-ups for the study twice a year, and will receive regular [health care](#) through their own health care providers.

The study will be double-blinded, so neither participants nor the study's clinical staff will know who is receiving vitamin D and who is receiving placebo. The study will continue until enough people have developed type 2 diabetes to be able to make a scientifically valid comparison between diabetes development in the two groups, likely about four years.

D2d builds on previous NIH-funded studies of methods to delay or prevent type 2 diabetes, including the Diabetes Prevention Program, which showed that, separately, lifestyle changes to lose a modest amount of weight and the drug metformin are both effective in slowing

development of type 2 diabetes in people with prediabetes. However, additional safe and effective preventative strategies are needed to stem the increasing numbers of people developing type 2 diabetes.

Provided by National Institute of Diabetes and Digestive and Kidney Diseases

Citation: Large study to examine if vitamin D prevents diabetes (2013, October 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-10-large-vitamin-d-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.