

## Synthetic vitamin D receptor ligands reduce murine kidney fibrosis

October 25 2013

Vitamin D deficiency has been associated with kidney disease including fibrosis. Some studies have even suggested that treatment with vitamin D or vitamin D analogs can reduce renal fibrosis; however, the pathways targeted by vitamin D therapy are not completely understood.

In this issue of the *Journal of Clinical Investigation*, Junn Yanagisawa and colleagues at the University of Tsukuba found that <u>vitamin</u> D binding to its receptor inhibited the TGF- $\beta$ /SMAD signaling pathway and prevented renal fibrosis in mice.

The authors then generated a synthetic ligand of the vitamin D receptor that, like vitamin D, reduced renal fibrosis; however, unlike vitamin D, this synthetic ligand did not promote hypercalcemia.

In the accompanying commentary Joseph Bonventre suggests that synthetic ligands of the vitamin D receptor should be further studied as therapeutics for patients with fibrotic diseases.

**More information:** A nonclassical vitamin D receptor pathway suppresses renal fibrosis, *J Clin Invest*. <u>DOI: 10.1172/JCI67804</u> Antifibrotic vitamin D analogs, *J Clin Invest*. 2013;123(11):4570–4573. <u>DOI: 10.1172/JCI72748</u>

Provided by Journal of Clinical Investigation



Citation: Synthetic vitamin D receptor ligands reduce murine kidney fibrosis (2013, October 25) retrieved 5 May 2024 from https://medicalxpress.com/news/2013-10-synthetic-vitamin-d-receptor-ligands.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.