

# Researchers find acute kidney injury predicts poor outcomes for dialysis patients

November 7 2013

---

Two University of Cincinnati (UC) researchers, in collaboration with other investigators, have found that patients who suffered from acute kidney injury (AKI) in the two-year period prior to going on dialysis were 1.5 times as likely to die in their first year of dialysis compared to those patients without AKI.

The findings will be presented Thursday, Nov. 7, at the American Society of Nephrology's Kidney Week in Atlanta.

Charuhas Thakar, MD, associate professor in the division of nephrology and hypertension, and Anthony Leonard, PhD, biostatistician and research assistant professor of family and community medicine, teamed up with Timmy Lee, MD, University of Alabama at Birmingham, and Pratik Parikh, PhD, Wright State University in Dayton, Ohio, to conduct this research.

The team examined the medical records of 47,327 [patients](#) in the United States Renal Data System who started [dialysis](#) from Jan. 1 to Dec. 31, 2008, and had Medicare data available in the preceding two years, and followed their mortality rates until Dec. 31, 2010.

"At present, when we start patients on chronic dialysis, we do not necessarily consider AKI episodes in the recent history to be a prognostic indicator," Thakar says. "Many times this information is not even available. Our findings suggest, however, that AKI occurrence prior to chronic dialysis initiation should alert both the patient and the

physician regarding a potentially poor outcome in the first year of dialysis."

AKI is marked by a sudden, temporary and sometimes fatal loss of [kidney](#) function. Rising incidence of this condition is related to an aging U.S. population and increasing need for hospitalization, Thakar says.

Thakar says the causes of AKI related to hospitalization include low blood pressure, dehydration, sepsis, heart failure and major surgeries such a cardiovascular surgery.

Researchers also found that AKI during the pre-dialysis period reduces the likelihood that patients will start dialysis with a functioning arterio-venous access, which provides the safest level of [dialysis care](#) for patients, Thakar explains.

"This finding indicates that AKI as a facilitator of progression of [chronic kidney disease](#) may impact the optimal preparation for dialysis leading to lack of preferred venous access for dialysis," Thakar adds.

About 400,000 people are on some form of dialysis in the United States, according to the National Kidney and Urologic Diseases Information Clearinghouse (NKUDIC).

Despite advances in technology and medical care these patients experience a high rate of mortality and reduced quality of life, Thakar says. "Recent studies have already established a link between the occurrence of AKI and growth of [chronic dialysis](#) population," Thakar explains. "Findings from our analysis add a novel dimension to this field by indicating that AKI may have long-lasting effects on dialysis outcomes as well."

Thakar says physicians should take greater care in reducing the exposure

of patients to kidney toxic medicines, ensuring early detection and treatment of sepsis and taking proper precautions during the perioperative period as some of the "common sense" steps to minimize the risk of AKI.

Thakar says the study has its limitations, adding: "These relationships are observational and not causally linked. More research needs to be done to understand whether this association plays out due to impacts on processes of care, or biology or both."

Thakar says that AKI may also be a marker of severity of illness in patients with dialysis and that further investigation is needed.

Provided by University of Cincinnati Academic Health Center

Citation: Researchers find acute kidney injury predicts poor outcomes for dialysis patients (2013, November 7) retrieved 10 April 2024 from <https://medicalxpress.com/news/2013-11-acute-kidney-injury-poor-outcomes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--