


Consumers order a less unhealthy meal when the menu has nutritional labeling

November 18 2013

Calories Purchased for a Single Meal



On average, customers at full-service restaurants purchased food that had approximately 1,600 food calories (kcal) -- a total that rose to 1,800 calories when also counting beverages. Most people need only 2,000 calories for an entire

day, so a single meal approximated a full day's worth of calories. Purchased meals had an average sodium content of 3,200 milligrams, with an average of 35 grams of saturated fat. These numbers far exceed recommended limits for an entire day. For most people, recommended daily limits are 2,300 milligrams sodium and 20 grams of saturated fat. Credit: Drexel University

An evaluation team led by the Drexel University School of Public Health has published a new study demonstrating that customers of full-service restaurants use nutritional labeling on menus to make healthier food choices.

"This is the first field-based study of mandatory [menu labeling](#) laws that found a large overall adjusted difference in [calories](#) between customers who dined at labeled restaurants when compared to unlabeled restaurants—about 155 fewer calories purchased," said Amy Auchincloss, PhD, an assistant professor in the Drexel University School of Public Health and lead author of the study.

Overall, customers at restaurants with menu labels purchased food with 151 fewer calories (155 fewer calories when counting beverages), 224 milligrams less sodium and 3.7 grams less saturated fat compared to customers at restaurants without menu labels.

Almost 80 percent of customers at labeled restaurants reported seeing labels, and 26 percent of all customers reported using them when deciding what to order. The customers who reported they used labels purchased 400 fewer calories (representing a relative difference of 20 percent), 370 milligrams less sodium and 10 grams less saturated fat than the overall average.

Nevertheless, even consumers who used the labels purchased oversized

meals that, on average, far exceeded what could be considered "healthy" – highlighting the difficulty for consumers when dining out. The authors argue for a need to do more to help consumers to eat sensibly and to encourage portion control, among other findings published online today ahead of print in the December issue of the *American Journal of Preventive Medicine*.

Americans currently purchase at least a third of the calories they consume on food prepared away from home – so providing detailed nutritional information on menus and on packaged foods is a commonly touted tactic to educate consumers and encourage them to make healthier choices.

"While previous studies have shown mixed impacts of menu labeling in fast food settings, this study suggests that nutrition information may be particularly useful in full-service restaurants," said Donald F. Schwarz, MD, health commissioner for the City of Philadelphia and a co-author of the study.

Philadelphia's menu labeling law requires full-service chain restaurants with more than 15 locations nationwide to list values for calories, sodium, fat and carbohydrates for each item on all printed menus. Fast food restaurants must list calories on their menus boards and make the other nutrition information available upon request. Philadelphia's law is unique in requiring more than just calories on menus. The enactment of this law in 2010 created the opportunity to observe whether menu labeling affects what consumers purchase – by comparing what happens at multiple locations of a single full-service chain restaurant, within and outside of city limits.

Menu labeling will expand nationwide when the Patient Protection and Affordable Care Act is fully implemented; at that time, all fast-food and full-service restaurant chains with more than 20 locations will be

required to provide [nutrition information](#) at the point of purchase.

In this study, Auchincloss and colleagues from Drexel, the Philadelphia Department of Public Health and University of Pennsylvania, assessed whether food purchases at full-service restaurants varied depending on the presence of labeling. They collected 648 customer surveys and transaction receipts at seven restaurant outlets of one large full-service restaurant chain. Two outlets had menu labeling, while five outlets did not. The authors looked at differences in calories and nutrients purchased between those who dined at outlets with menu labeling and those who did not, and at customers' reported use of nutritional information when ordering.

On average, customers purchased food that had approximately 1,600 food calories (kcal) – a total that rose to 1,800 calories when also counting beverages. Most people need only 2,000 calories for an entire day, so a single meal approximated a full day's worth of calories. Purchased meals had an average sodium content of 3,200 milligrams, with an average of 35 grams of saturated fat. These numbers far exceed recommended limits for an entire day. For most people, recommended daily limits are 2,300 milligrams sodium and 20 grams of saturated fat.

"When you compare the average intake with the recommended daily intake, these consumers purchased almost all their calories, and more than the recommended sodium and saturated fat in just one meal," said Beth Leonberg, an assistant clinical professor and director of the didactic program in dietetics in Drexel's College of Nursing and Health Professions, who co-authored the study. "In order to not exceed recommended intakes for the day, most adults should consume fewer than 750 calories, 750 milligrams of sodium and 8 grams of [saturated fat](#) in a single meal."

The authors concluded that current efforts don't go far enough to help

consumers to eat sensibly and to encourage portion control. Educating consumers about menu labeling may further increase the small observed impact on healthier consumer choices.

"We also need to pursue approaches that make the healthy choice the default," said Giridhar Mallya, MD, director of policy and planning for the Philadelphia Department of Public Health and a co-author of the study. "This might include product reformulation, promoting healthier options on menus, and offering smaller portion sizes."

More information: [dx.doi.org/10.1016/j.amepre.2013.07.014](https://doi.org/10.1016/j.amepre.2013.07.014)

Provided by Drexel University

Citation: Consumers order a less unhealthy meal when the menu has nutritional labeling (2013, November 18) retrieved 23 April 2024 from <https://medicalxpress.com/news/2013-11-consumers-unhealthy-meal-menu-nutritional.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.