

New research may improve early detection of dementia

November 12 2013

Using scores obtained from cognitive tests, Johns Hopkins researchers think they have developed a model that could help determine whether memory loss in older adults is benign or a stop on the way to Alzheimer's disease.

The risk of developing [dementia](#) increases markedly when a person is diagnosed with [mild cognitive impairment](#), a noticeable and measurable decline in intellectual abilities that does not seriously interfere with daily life. But physicians have no reliable way to predict which people with mild cognitive impairment are likely to be in the 5 to 10 percent a year who progress to dementia.

In a proof-of-concept study, the Johns Hopkins investigators analyzed records of 528 people age 60 and over, who were referred to the Johns Hopkins Medical Psychology Clinic for cognitive testing as part of a dementia work-up between 1996 and 2004. The results were compared to those of 135 healthy older adults who participated in a study of normal aging. Both groups completed tests of memory, language, attention, processing speed and drawing abilities from which 13 scores were recorded.

Since each person is naturally more skillful in some areas than in others, the scores of healthy adults showed a symmetrical, bell-shaped range: Most of their scores were high, a few were a bit lower, and a few were even lower. By grouping the patients into cohorts based on the severity of their dementia, the researchers found a trend in the [test scores](#) that is

likely to mimic the deterioration of an individual's scores over time.

At the outset, he says, Alzheimer's disease subtly disrupts some mental abilities, while leaving others intact. Thus, well before a person develops clear cognitive impairment, his or her performance declines slightly on a few measures. When shown on a graph, these changes cause the healthy symmetric, bell-shaped curve to shift and become asymmetrical.

Regardless of how low a person's test scores were, the researchers determined that lopsidedness in their score distribution correlated with dementia. They predicted that people with low scores that were evenly distributed were not likely to develop dementia. But those with clearly lopsided test score distributions on the 13 measures administered were already experiencing varying levels of dementia.

"Departures from the normal bell-shaped pattern of variability on [cognitive tests](#) might determine which people with low scores develop dementia," says David J. Schretlen, Ph.D., a professor of psychiatry and behavioral sciences at the Johns Hopkins University School of Medicine and leader of a study published online Nov. 12 in the journal *Neuropsychology*.

Since these declines can be subtle, the researchers also increased the precision of cognitive testing by accounting for the effects of age, sex, race and education on test performance.

The challenge for doctors, Schretlen explains, is that most normal, healthy people will produce a few low scores on cognitive testing. That makes it nearly impossible to know at the outset whether a patient who reports forgetfulness and produces one or two low scores has a benign form of mild cognitive impairment, or is in the earliest stage of dementia. As a result, doctors often tell such patients to return for follow-up testing in a year or two.

But if future research confirms it, this new statistical model could help doctors get the prognosis right earlier in the disease, at the first visit, and start treating patients accordingly.

Mostly, Schretlen says, doctors could use the new model to reassure patients who are not at risk of dementia, while fast-tracking interventions for those who are. Because there currently are no effective treatments for Alzheimer's disease, those likely headed that way could be counseled to take the good time they have to organize their affairs, and do things they have always wanted to do. They also could be fast-tracked into clinical trials of medications to slow the progression of dementia.

"If we are going to have any hope of helping patients with Alzheimer's disease, we need to do it as early as possible," Schretlen says. "Once the brain deteriorates, there's no coming back."

Recent failures of drugs in late-stage clinical trials for Alzheimer's disease have been a real blow, he adds, but new treatments are being developed.

The new way of reading existing test scores follows a 2008 study by the same researchers showing that one of every six healthy adults scored poorly on two or more of 10 tests in a brief cognitive battery—even though there was nothing wrong with them.

The main reason it is difficult to tell whether older people have benign mild cognitive impairment or not is because they are not routinely screened for [cognitive impairment](#), he says. A visit to a specialist comes only after someone has noticed symptoms, and then cognitive testing is interpreted without the benefit of a baseline assessment. What would solve this problem, he says, would be for everyone over the age of 55 to get routine neurocognitive testing every five years.

Provided by Johns Hopkins University School of Medicine

Citation: New research may improve early detection of dementia (2013, November 12) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-11-early-dementia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.