

Researchers identify a histone demethylase associated with non-small cell lung cancer

November 8 2013

Non-small cell lung cancer (NSCLC) is one of the leading causes of cancer deaths worldwide. Current therapies that target cellular kinases have been effective for some patients: however, many individuals with NSCLS do not respond. Recent studies of other cancers indicate that the global changes in DNA methylation patterns may drive tumor formation.

In this issue of the *Journal of Clinical Investigation*, MinGyu Lee and colleagues at MD Anderson Cancer Center evaluated histone methylation modifications in NSCLS cell lines and determined that the histone demethylase KDM2A was upregulated in NSCLC cell lines. Furthermore, induced KDM2A expression increased <u>cell proliferation</u> and invasiveness through activation of the ERK1/2 signaling pathway. Evaluation of tumors from NSCLS patients uncovered a correlation between increased levels of KDM2A and poor prognosis.

This study suggests that therapies targeting KDM2A may benefit a subset of NSCLS patients.

More information: KDM2A promotes lung tumorigenesis by epigenetically enhancing ERK1/2 signaling, *J Clin Invest*. DOI: 10.1172/JCI68642

Provided by Journal of Clinical Investigation



Citation: Researchers identify a histone demethylase associated with non-small cell lung cancer (2013, November 8) retrieved 5 May 2024 from https://medicalxpress.com/news/2013-11-histone-demethylase-non-small-cell-lung.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.