

Researchers link lack of sleep in teens to higher risk of illness

November 14 2013

Newly released findings from Bradley Hospital published in the *Journal of Sleep Research* have found that acute illnesses, such as colds, flu, and gastroenteritis were more common among healthy adolescents who got less sleep at night. Additionally, the regularity of teens' sleep schedules was found to impact their health. The study, titled "Sleep patterns are associated with common illness in adolescents," was led by Kathryn Orzech, Ph.D. of the Bradley Hospital Sleep Research Laboratory.

Orzech and her team compared three outcomes between longer and shorter sleepers: number of illness bouts, illness duration, and school absences related to illness. The team found that bouts of illness declined with longer sleep for both male and female high school students. Longer sleep was also generally protective against school absences that students attributed to illness. There were gender differences as well, with males reporting fewer illness bouts than females, even with similar sleep durations.

Orzech's team analyzed total sleep time in teens for six-day windows both before and after a reported illness and found a trend in the data toward shorter sleep before illness vs. wellness. Due to the difficulty of finding teens whose illnesses were spaced in such a way to be statistically analyzed, Orzech also conducted qualitative analysis, examining individual interview data for two short-sleeping males who reported very different illness profiles. This analysis suggested that more irregular sleep timing across weeknights and weekends (very little sleep during the week and "catching up" on sleep during the weekend), and a preference

for scheduling work and social time later in the evening hours can both contribute to differences in illness outcomes, conclusions that are also supported in the broader adolescent sleep literature.

"Some news reaches the general public about the long-term consequences of [sleep deprivation](#), such as the links between less sleep and weight gain," said Orzech. "However, most of the studies of sleep and health have been done under laboratory conditions that cannot replicate the complexities of life in the real world. Our study looked at rigorously collected sleep and illness data among adolescents who were living their normal lives and going to school across a school term."

"We showed that there are short-term outcomes, like more acute [illness](#) among shorter-sleeping adolescents, that don't require waiting months, years or decades to show up," Orzech continued. "Yes, poor sleep is linked to increased cardiovascular disease, to high cholesterol, to obesity, to depression, etc., but for a teenager, staying healthy for the dance next week, or the game on Thursday, may be more important. This message from this study is clear: Sleep more, and more regularly, get sick less."

Mary Carskadon, Ph.D., director of the Bradley Hospital Sleep Research Laboratory, commented on Orzech's study, "We have long been examining the sleep cycles of teenagers and how we might be able to help adolescents - especially high school students - be better rested and more functional in a period of their lives where sleep seems to be a luxury." Carskadon continued, "In the future, these findings identifying specific issues in individual [sleep patterns](#) may be a useful way to help adolescents begin to prioritize [sleep](#)."

Provided by Lifespan

Citation: Researchers link lack of sleep in teens to higher risk of illness (2013, November 14)

retrieved 6 May 2024 from

<https://medicalxpress.com/news/2013-11-link-lack-teens-higher-illness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.