

Study finds few patients with newly-diagnosed hyperlipidemia receive recommended thyroid screening

November 13 2013

Despite current guidelines that recommend newly diagnosed high-cholesterol patients have a TSH blood test done to make sure they do not have hypothyroidism, researchers from Boston University School of Medicine (BUSM) and Boston Medical Center (BMC) have found that only about half of these patients were screened for thyroid dysfunction. The findings, which appear online in *JAMA Internal Medicine*, show the current guidelines may be underutilized.

Hypothyroidism (an underactive thyroid) is a relatively common cause of high cholesterol. Between one and 13 percent of [patients](#) with high cholesterol are thought to have hypothyroidism. Current guidelines recommend that patients newly diagnosed with high cholesterol should have a TSH blood test done to make sure they do not have hypothyroidism.

Researchers reviewed the records of adult patients with a new diagnosis of high cholesterol between 2003-2011 and found 8,795 patients with new high cholesterol diagnoses within the study period. Thyroid function was checked within six months of the high cholesterol diagnosis in 49.5 percent of the patients. Of those, 226 (5.2 percent) had an elevated TSH level consistent with hypothyroidism. Among the patients with hypothyroidism, 114 (50.7 percent) were treated with [thyroid hormone](#). Only 25 percent of the patients treated with thyroid hormone (versus 44 percent of the patients not treated with thyroid hormone) were also

prescribed a lipid-lowering medication within one year.

"We performed this retrospective study to see how many patients with newly diagnosed high cholesterol were having their thyroid function checked and found that only about half of these patients were screened for [thyroid dysfunction](#), despite current guidelines," explained corresponding author Elizabeth Pearce, MD, associate professor of medicine at BUSM and endocrinologist at BMC. "We conclude that the low rate of thyroid function testing in patients with new-onset hyperlipidemia demonstrates the need for greater awareness of current guidelines."

According to Pearce, future studies are needed to better understand reasons for low [thyroid function](#) screening rates among patients with high cholesterol, and the cost-effectiveness of [hypothyroidism](#) screening and treatment for patients with [high cholesterol](#).

Provided by Boston University Medical Center

Citation: Study finds few patients with newly-diagnosed hyperlipidemia receive recommended thyroid screening (2013, November 13) retrieved 18 April 2024 from <https://medicalxpress.com/news/2013-11-patients-newly-diagnosed-hyperlipidemia-thyroid-screening.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--