

## **Results of the REPRISE II trial reported**

November 1 2013

In a clinical trial, a second-generation transcatheter aortic valve demonstrated low rates of complications that are sometimes seen in transcatheter aortic valve replacement (TAVR), including challenges with positioning, post-procedure paravalvular aortic regurgitation, vascular complications, and stroke.

The findings were presented today at the 25th annual Transcatheter Cardiovascular Therapeutics (TCT) scientific symposium. Sponsored by the Cardiovascular Research Foundation (CRF), TCT is the world's premier educational meeting specializing in interventional cardiovascular medicine.

The valve studied in REPRISE II is fully retrievable and repositionable with a unique adaptive seal intended to minimize paravalvular regurgitation, a complication that has been associated with higher mortality among patients undergoing TAVR. In this prospective, single arm, multicenter study, symptomatic patients at high risk for surgery received the Lotus valve to treat calcific aortic stenosis.

The trial enrolled 120 patients; mean age was  $84.4\pm5.3$  years, 56.7 percent were female, and 75.8 percent were considered New York Heart Association (NYHA) Class III or IV. The mean Society of Thoracic Surgeons score was  $7.1\pm4.6$  percent and all patients were confirmed by their site heart team to be at high risk for surgery due to frailty or associated comorbidities.

The valve was successfully implanted in all 120 patients with valve



repositioning and retrieval performed as needed. There was no embolization, ectopic valve deployment, or need for implantation of a second prosthetic valve.

The primary device performance endpoint was the mean aortic valve pressure gradient at 30 days compared to a performance goal of 18 mmHg; the primary safety endpoint was 30-day mortality. The primary device performance endpoint was met with a 30 day mean <u>aortic valve</u> pressure gradient of  $11.5\pm5.2$  mmHg; mean effective orifice area was  $1.7\pm0.4$  cm2. All cause mortality and disabling stroke were low at 30 days (4.2 percent and 1.7 percent, respectively). Additional clinical event rates were consistent with those reported for other valves. Aortic regurgitation at 30 days was negligible in 99.0 percent of <u>patients</u> (78.3 percent none, 5.2 percent trace, and 15.5 percent mild).

"These findings suggest this valve, which is a differentiated, second generation TAVR device, will be a valuable addition for the treatment of severe <u>aortic stenosis</u>," said the lead investigator, Ian T. Meredith, MBBS, PhD. Dr. Meredith is Director of Monash HEART and the Executive Director of the Monash Cardiovascular Research Centre. He is also a Professor of Medicine at Monash University in Melbourne, Australia.

Provided by Cardiovascular Research Foundation

Citation: Results of the REPRISE II trial reported (2013, November 1) retrieved 3 June 2024 from <u>https://medicalxpress.com/news/2013-11-results-reprise-ii-trial.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.