

Researchers discover promising new treatment to help people with spine injuries walk better

November 27 2013

Scientists may have found a new treatment that can help people with spinal cord injuries walk better. The research is published in the November 27, 2013, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

"About 59 percent of all <u>spinal injuries</u> are incomplete, leaving pathways that could allow the <u>spinal cord</u> to change in a way that allows people to walk again. Unfortunately, usually a person affected by this type of spinal injury seldom recovers the ability to walk normally," said study author Randy D. Trumbower, PT, PhD, with Emory University in Atlanta. "Our research proposes a promising new way for the spinal cord to make the connections needed to walk better."

The research involved 19 people with spine injuries between levels C2 and T12, no joint shortening, some controlled ankle, knee, and hip movements, and the ability to walk at least one step without human assistance. Research team members were based at Emory University, Georgia Institute of Technology and Shepherd Center in Atlanta, the Rehabilitation Institute of Chicago and the University of Wisconsin, Madison.

The participants were exposed to short periods of breathing low <u>oxygen</u> <u>levels</u>, which is called hypoxia. The participants breathed through a mask for about 40 minutes a day for five days, receiving 90-second periods of



low oxygen levels followed by 60 seconds of normal oxygen levels. The participants' walking speed and endurance was tested before the study started, on the first and fifth days of <u>treatment</u>, and again one and two weeks after the treatment ended.

The participants were divided into two groups. In one, nine people received either the treatment or a sham treatment where they received only normal oxygen levels. Then two weeks later they received the other treatment. In the other group, the participants received the treatment or sham treatment and then were asked to walk as fast as they could for 30 minutes within one hour of the treatment, then received the other treatment two weeks later.

Those who received just the hypoxia treatment increased their walking speed on a test of walking 10 meters, walking an average of 3.8 seconds faster than when they did not receive the treatment.

Those who had the treatment plus walking increased their endurance on a test of how far they could walk in six minutes by an average of 100 meters, which was more than a 250-percent increase compared to those who had the sham treatment plus walking.

All participants improved their ability to walk. More than 30 percent of all <u>participants</u> increased their <u>walking speed</u> by at least a tenth of a meter per second and more than 70 percent increased their endurance by at least 50 meters.

"One question this research brings to light is how a treatment that requires people to take in low levels of oxygen can help movement, let alone in those with compromised lung function and motor abilities," said Michael G. Fehlings, MD, PhD, with the University of Toronto in Canada, who wrote a corresponding editorial on the study. "A possible answer is that spinal serotonin, a neurotransmitter, sets off a cascade of



changes in proteins that help restore connections in the spine."

Trumbower cautions that chronic or sustained hypoxia in untrained hands may cause serious injury and should not be attempted outside the scope of a supervised medical treatment.

Provided by American Academy of Neurology

Citation: Researchers discover promising new treatment to help people with spine injuries walk better (2013, November 27) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2013-11-treatment-people-spine-injuries.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.