

Angioplasty possibly no better than drug therapy

December 9 2013

For patients with stable coronary artery disease (CAD) who are not experiencing a heart attack and an abnormal stress test, treatment of their narrowed arteries by the common procedure of angioplasty may not provide additional benefits compared to drug therapy alone. This finding results from a survey of more than 4,000 patients with myocardial ischemia, or inadequate circulation, led by cardiologists at Stony Brook University School of Medicine. The survey results are published in the online first edition of *JAMA Internal Medicine*.

In the research paper titled "Percutaneous Coronary Intervention Outcomes in Patients With Stable Obstructive Coronary Artery Disease and Myocardial Ischemia: A Collaborative Meta-analysis of Contemporary Randomized Clinical Trials," David L. Brown, MD, and Kathleen Stergiopoulos, MD, PhD, Professors in the Department of Medicine, Division of Cardiovascular Medicine at Stony Brook University School of Medicine, worked with colleagues internationally to design a data study that combined data from clinical trials performed between 1970 and 2012 of [patients](#) who had either [percutaneous coronary intervention](#) (PCI), or angioplasty, plus [drug therapy](#), or drug therapy alone to treat their CAD.

Each of the clinical studies within the analysis reported outcomes of death and nonfatal [myocardial infarction](#) reported. Additionally, to reflect contemporary medical and interventional practice, inclusion criteria required stent implantation in at least 50 percent of the PCI procedures and statin medications to lower cholesterol in at least 50

percent of patients in both the PCI and drug therapy alone groups. This led to a total of five clinical trials yielding 4,064 patients with [myocardial ischemia](#) diagnosed by exercise stress testing, nuclear or echocardiographic stress imaging, or fractional flow reserve.

The researchers reviewed outcomes data up to five years post PCI or drug treatment alone. They analyzed all-cause death, non-fatal myocardial infarction, unplanned revascularization, and angina in the patients.

The analysis showed all-cause death rates between the two groups was not significantly different – 6.5 percent for patients receiving PCI and drug therapy versus 7.3 percent for patients receiving drug therapy alone. There was little difference in the rates of non-fatal myocardial infarction (9.2 percent with PCI vs. 7.6 percent drug therapy) and recurrent or persistent angina (20.3 percent vs. 23.3). The rate of unplanned revascularization was slightly different but not statistically significant (18.3 percent vs. 28.4 percent).

"If our findings are confirmed in ongoing trials, many of the more than 10 million stress tests performed annually and subsequent revascularizations may be unnecessary," said Dr. Brown.

He cautioned that additional studies beyond data analyses of [clinical trials](#) are necessary to fully determine if practices with PCI in stable CAD patients needs to be re-evaluated, and if so, under what circumstances and in which patient populations.

More information: Stergiopoulos K, Boden WE, Hartigan P, et al. Percutaneous Coronary Intervention Outcomes in Patients With Stable Obstructive Coronary Artery Disease and Myocardial Ischemia: A Collaborative Meta-analysis of Contemporary Randomized Clinical Trials. *JAMA Intern Med.* 2013. [DOI:](#)

[10.1001/jamainternmed.2013.12855](https://doi.org/10.1001/jamainternmed.2013.12855)

Provided by Stony Brook University

Citation: Angioplasty possibly no better than drug therapy (2013, December 9) retrieved 5 May 2024 from <https://medicalxpress.com/news/2013-12-angioplasty-possibly-drug-therapy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.