

BMI not linked to pain after exercise rehab for back pain

December 12 2013



(HealthDay)—For individuals with chronic low back pain (cLBP), body mass index (BMI) is not significantly associated with self-reported pain and disability, according to a study published in the Dec. 1 issue of *Spine*.

Cristy Brooks, from the University of Western Sydney, and colleagues retrospectively analyzed changes in self-reported pain and disability among 128 males and females with cLBP who performed eight weeks of exercise, consisting of three to five exercise sessions (minimum of one supervised session) per week.

The researchers found that there were no baseline relationships between BMI and self-reported pain and disability ($P = 0.349$ and 0.314 , respectively). No association was seen for exercise-related changes in

pain and disability with BMI at baseline ($P = 0.938$ and 0.873 , respectively) or with changes in BMI ($P = 0.402$ and 0.854 , respectively). Furthermore, exercise-based pain and disability changes were not predicted by BMI.

"The reliance on BMI as a sole measurement of obesity in cLBP research may be unwarranted," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: BMI not linked to pain after exercise rehab for back pain (2013, December 12)
retrieved 20 April 2024 from
<https://medicalxpress.com/news/2013-12-bmi-linked-pain-rehab.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--