

Five-minute intervention can help patients with substance and behavioral problems

December 10 2013, by Helen Dodson



A brief 5 to 10 minute intervention in an emergency department can provide effective help to patients with substance and behavioral problems, and should be broadly implemented, say the authors of a new Yale study. Their recommendations appear in the December issue of *Health Affairs*.

Millions of patients seek treatment in emergency departments (ED) each year for substance use and [behavioral disorders](#). Hospitals and primary care facilities have limited capacity to treat these patients, who often are

poor, uninsured, and sick.

Adult patients who enter the ED are more likely to smoke, drink, use illicit substances, and misuse prescription drugs than are members of the general population, the authors write. A growing body of evidence suggests that screening, providing a brief intervention, and referring these patients to treatment (a protocol known as SBIRT) can be an effective way to help them. SBIRT can reduce these harmful habits among ED patients, the authors write, having already proven effective in helping people quit smoking and reduce alcohol and marijuana use.

SBIRT is an evidence-based technique that incorporates principles of motivational interviewing, empathy, positive framing, reflective listening, and gentle education to encourage people to change risky behaviors. In addition to being an effective counseling technique, say the researchers, there is evidence that investing in SBIRT can result in healthcare cost savings by reducing the use of the ED.

The authors are Steven Bernstein, M.D., professor of [emergency medicine](#) at Yale School of Medicine, and Dr. Gail D'Onofrio, professor and chair of emergency medicine at Yale School of Medicine.

More information: Bernstein, Steven L. "A Promising Approach For Emergency Departments To Care For Patients With Substance Use And Behavioral Disorders." *Health Aff* , December 2013 32:122122-2128; [DOI: 10.1377/hlthaff.2013.0664](https://doi.org/10.1377/hlthaff.2013.0664)

Provided by Yale University

Citation: Five-minute intervention can help patients with substance and behavioral problems (2013, December 10) retrieved 20 March 2024 from

<https://medicalxpress.com/news/2013-12-five-minute-intervention-patients-substance-behavioral.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.