

Greater dietary fiber intake associated with lower risk of heart disease

December 19 2013

Greater dietary fibre intake is associated with a lower risk of both cardiovascular disease and coronary heart disease, a paper published today in *BMJ* suggests.

In recent years, a decline in both cardiovascular disease (CVD) and coronary [heart disease](#) (CHD) has been seen in some European countries and the United States. However, it still remains a significant issue accounting for almost half (48%) and a third (34%) of all deaths in Europe and the United States.

Many studies have examined the relationship between [dietary fibre](#) or fibre-rich foods and CVD risk factors.

Researchers at the University of Leeds reviewed literature published since 1990 in healthy populations concerning dietary fibre intake and CVD risk. They took data from six electronic databases. Cohorts of data were used from the US, Europe, Japan and Australia.

They looked at the following fibre intake: total, insoluble (whole grains, potato skins etc), soluble (legumes, nuts, oats, barley etc), cereal, fruit, vegetable and other sources.

Results from analyses of total, insoluble, fruit and vegetable fibre intake showed that the likelihood of a CVD or CHD event steadily lowers with increasing intake.

In soluble fibre, a higher reduction was seen in CVD risk than CHD risk and for cereal fibre, the reduced risk of CHD was stronger than the association with CVD.

A significantly lower risk of both CVD and CHD was observed with every additional 7g per day of fibre consumed. The researchers say these findings are aligned with current recommendations to increase fibre intake and demonstrate a large risk reduction with an achievable increase in daily fibre intake and say this could "potentially impact on many thousands of individuals."

They add that an additional 7g of fibre can be achieved through one portion of wholegrains (found in bread, cereal, rice, pasta) plus a portion of beans / lentils or two to four servings of fruit and vegetables.

The researchers conclude that "diets high in fibre, specifically from cereal or vegetable sources ... are significantly associated with lower risk of CHD and CVD and reflect recommendations to increase intake." Greater intake from fruit fibre was associated with lower CVD risk. They recommend further work on the association with soluble or insoluble types of fibre.

In an accompanying editorial, Dr Robert Baron, Professor of Medicine at the University of California, says this study "increases our confidence that benefit, as reflected by reduced [cardiovascular disease](#) and [coronary heart disease](#) events, will in fact accrue with higher dietary fibre intakes."

He says that teaching patients to eat whole grains is still challenging, but that encouraging the increase of fibre gradually as well as drinking adequate amounts of water are other practical recommendations. The recommendation to consume diets with adequate amounts of dietary fibre "may turn out to be the most important nutrition recommendation

of them all," he concludes.

More information: www.bmj.com/cgi/doi/10.1136/bmj.f6879
www.bmj.com/cgi/doi/10.1136/bmj.f7401

Provided by British Medical Journal

Citation: Greater dietary fiber intake associated with lower risk of heart disease (2013, December 19) retrieved 4 May 2024 from <https://medicalxpress.com/news/2013-12-greater-dietary-fiber-intake-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--