


Inadequate pregnancy weight gain a risk factor for infant mortality

December 19 2013



Study findings show that women who gain too little weight during pregnancy are more likely to give birth to babies who die in infancy. In order to improve the likelihood of healthy outcomes for mothers and infants, healthcare professionals should provide childbearing women with gestational weight gain goals that are specific to their individual BMI. Credit: University of Maryland

Women who do not gain enough weight during pregnancy are at increased risk of losing their baby in its first year of life, according to a

new study by researchers in the University of Maryland School of Public Health (UMD SPH). This study examined the relationship between gestational weight gain, mothers' body mass index (BMI) before and during pregnancy, and infant mortality rates. One-quarter of the more than 159,000 women in the study gained too little weight during pregnancy, and these mothers were more likely to give birth to babies who died in infancy than the women who gained a normal or even excessive amount of weight during pregnancy.

The study is published in the *American Journal of Public Health*. It was conducted by Dr. Regina Davis, associate executive director of the American Public Health Association, Dr. Sandra Hofferth, professor, and Dr. Edmond Shenassa, associate professor. All three are affiliated with the UMD SPH Maternal and Child Health Program, Department of Family Science, and Hofferth and Shenassa are faculty associates of the Maryland Population Research Center.

"Our study showed that gaining too little [weight](#) during [pregnancy](#) is a risk factor for [infant mortality](#) for all but the heaviest women," explained Dr. Davis. Gaining more weight than recommended was not a risk factor for infant mortality but may be related to subsequent maternal health problems. "It is important that childbearing women have pregnancy [weight gain](#) goals that are specific to their individual BMI," added Dr. Hofferth, the study's senior author.

According to guidelines published by the Institute of Medicine (IOM), underweight women should gain between 28 and 40 pounds during pregnancy, while normal weight women are expected to gain 25-35 pounds, overweight women 15-25 pounds, and obese women 11-20 pounds. Results from this study showed that only about a third of women gained an amount of weight that was within the recommended guidelines for their body size. Nearly 25% of the women in the study gained an inadequate amount of weight, while another 41% gained too much.

Davis, Hofferth, and Shenassa analyzed data collected from 159,244 mothers who gave birth to live, single babies between 2004 and 2008 in order to determine whether there was a link between gestational weight gain (GWG), mothers' [body mass index](#) (BMI), and infant mortality. The women in the study responded to telephone or written questionnaires within nine months after their babies' birth, and their information was recorded in the Pregnancy Risk Assessment Monitoring System (PRAMS) for use by future researchers. The study team analyzed the PRAMS data to learn whether mothers' BMI before and during pregnancy might alter the influence of GWG on infant mortality.

Infant mortality risks in the study sample were 3.9% among infants of mothers who gained an inadequate amount of weight during pregnancy, 1.2% among infants of mothers who gained an adequate amount of weight, and .7% among mothers who gained more than the recommended amount. Mothers' pre-pregnancy BMI was also a key factor in infant survival. Mothers who were underweight before pregnancy and gained too little weight during pregnancy had six times the normal rate of infant mortality. Even among overweight women, inadequate weight gain was associated with a two-fold elevation in the risk of mortality. Only children born to obese women were protected from the effects of inadequate weight gain. In contrast, gaining more than the recommended amount of weight was not associated with risk to the infant among mothers in any weight category. Obese mothers who gained an excessive amount of weight actually had a 49% reduced likelihood of infant death.

This study confirms that only about a third of women in the US gain weight during pregnancy that is within the recommended range. In order to improve the likelihood of healthy outcomes for [mothers](#) and infants, healthcare providers should provide childbearing women with gestational weight gain goals that are specific to their individual BMI. Pregnant [women](#) can help ensure the health of their infants by monitoring their

weight gain, eating an appropriate amount of healthy foods, and engaging in reasonable physical activity.

More information: Study results are published in the *American Journal of Public Health* on December 19, 2013.

Provided by University of Maryland

Citation: Inadequate pregnancy weight gain a risk factor for infant mortality (2013, December 19) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-12-inadequate-pregnancy-weight-gain-factor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.