

## Study links sleep to mood disturbance and poor quality of life in obese

December 4 2013

---

A new study shows that poor sleep quality is strongly associated with mood disturbance and lower quality of life among people with extreme obesity.

Results show that 74.8 percent of participants were poor sleepers, and their mean self-reported sleep duration was only six hours and 20 minutes. Fifty-two percent of study subjects were anxious, and 43 percent were depressed. After controlling for age, sex, hypertension, diabetes, and [obstructive sleep apnea](#), sleep quality and [daytime sleepiness](#) were significantly associated with [mood disturbance](#) and quality of life impairment.

"There was a clear association between the [sleep problems](#) such as short [sleep duration](#) and the psychological disorders and with quality of life," said Dr. G. Neil Thomas, lead supervisor, study methodology lead and reader in epidemiology at the Department of Public Health, Epidemiology and Biostatistics at the University of Birmingham in the United Kingdom. "These associations remained significant even after adjusting for a range of potential confounders."

The study involved 270 patients with a mean body mass index (BMI) of 47.0 kg/m<sup>2</sup> who were consecutively enrolled in a regional specialist weight management service. They had a mean age of 43 years. Sleep disturbance, daytime sleepiness, mood and quality of life were assessed using standardized questionnaires.

The study results appear in the December issue of the journal *Sleep*.

"This study emphasizes the need for physicians to conduct routine screenings for sleep problems among people with severe obesity," said American Academy of Sleep Medicine President Dr. M. Safwan Badr. "Improving [sleep quality](#) and quantity will provide a physical, mental and emotional boost for people who are making the difficult lifestyle changes involved in managing obesity."

According to the authors, the potential role of sleep in the health and well-being of individuals with [severe obesity](#) is underappreciated. Although the cross-sectional design of the study did not allow for an examination of causality, the results suggest that the early detection of disturbed sleep could prevent the potential development and perpetuation of psychological problems among people with [extreme obesity](#).

"Despite the very high levels of problems in these patients, those involved with their care usually don't ask about sleep problems and often pay little heed to the psychological issues underlying the obesity," said Thomas. "The focus is often on treating the obesity and its consequences, such as diet and exercise interventions, rather than addressing its underlying cause, which may be psychological in nature, such as an unhappy marriage or job stress."

According to the Centers for Disease Control and Prevention, 35.7 percent of U.S. adults are obese with a BMI of 30 or higher. The CDC estimates that the annual medical cost of obesity in the U.S. was \$147 billion in 2008 dollars.

**More information:** "The Complex Associations Among Sleep Quality, Anxiety-Depression, and Quality of Life in Patients with Extreme Obesity" *Sleep*, 2013.

Provided by American Academy of Sleep Medicine

Citation: Study links sleep to mood disturbance and poor quality of life in obese (2013, December 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-12-links-mood-disturbance-poor-quality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.