

# Many people with diabetes still lose vision, despite availability of vision-sparing treatment

December 19 2013

---

Despite recent advances in prevention and treatment of most vision loss attributed to diabetes, a new study shows that fewer than half of Americans with damage to their eyes from diabetes are aware of the link between the disease and visual impairment, and only six in 10 had their eyes fully examined in the year leading up to the study.

The research, described online on Dec. 19 in *JAMA Ophthalmology*, also found that nearly half of those with [diabetes](#) and [eye](#) damage had not visited a clinician charged with managing their disease in that same time period.

High blood sugar levels associated with poorly controlled diabetes can damage the small blood vessels in the retina, the light-sensitive tissue lining the back wall of the eye. This can cause [poor blood flow](#) or leaking of fluids from these blood vessels, which brings on swelling of the retina. The condition, known as [diabetic macular edema](#), can cripple central vision and the ability to see detailed images, such as the face of someone looking directly at you. Left untreated for a year or longer, macular edema can lead to permanent vision loss.

"As a nation, we are woefully inadequate as [health care providers](#) in explaining to our patients with diabetes that the condition can have a detrimental effect on their eyes," says study leader Neil M. Bressler, M.D., a professor of ophthalmology at the Johns Hopkins University

School of Medicine and chief of the retina division at the Johns Hopkins Wilmer Eye Institute. "The earlier we catch [diabetic eye disease](#), the greater the likelihood that we can help patients keep their good vision. Clearly, this research shows how far we have to go to educate people about this frequent and feared complication."

People with diabetes have at least a 10 percent risk of developing diabetic macular edema during their lifetime, and estimates suggest that close to 745,000 of them in the United States have swelling in the macula, the center portion of the retina.

Until recently, 15 percent of patients who developed the condition and were treated for it with the standard laser therapy still lost their vision. Now, Bressler says, drugs injected into the eye reduce the swelling and risk of vision loss to less than five percent. With treatment, moreover, half of patients find their vision improves, making prompt diagnosis critical.

For the study, the Johns Hopkins-led team of researchers used data collected between 2005 and 2008 from Americans enrolled in the National Health and Nutrition Examination Survey (NHANES). Among the 798 people over the age of 40 with a self-reported diagnosis of type 2 diabetes and who had retinal imaging done, 48 had diabetic macular edema and were asked in the survey whether a physician had told them about the link between diabetes and vision problems (44.7 percent were). They were also asked whether they had seen a health care provider about their diabetes in the previous year (46.7 percent had), and whether they had received an eye examination, including pupil dilation, in the previous year (59.7 percent had). Some 30 percent of the individuals with diabetic macular edema already had some type of [vision loss](#) related to the disease.

Bressler says some people fail to see eye doctors or diabetes educators

because they lack insurance. He adds that most of the problem is likely a lack of understanding about the risks, and most people probably aren't referred to eye care specialists who can quickly determine retinal vulnerability.

"We can prevent a lot of [vision](#) impairment or blindness if we can just get these people into the medical system," Bressler says. Now that the extent of the problem is known, Bressler says, strategies can be developed to address issues of patient education, access to specialists and costs.

Provided by Johns Hopkins University School of Medicine

Citation: Many people with diabetes still lose vision, despite availability of vision-sparing treatment (2013, December 19) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-12-people-diabetes-vision-availability-vision-sparing.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--