

Assessing the US population's sodium intake

December 6 2013, by Rosalie Marion Bliss



An ARS food-survey tool proves accurate in estimating the amount of sodium consumed by volunteers. Credit: Peggy Greb

U.S. Department of Agriculture (USDA) scientists used an automated dietary survey tool they developed to accurately estimate how much sodium volunteers consumed as part of their daily diets.

The scientists, with USDA's Agricultural Research Service (ARS), found that the volunteers' [sodium intake](#) estimates were 90 to 93 percent accurate among men and women. Sodium intake has become a hot topic

as public policymakers address regulatory proposals aimed at lowering [sodium](#) in foods. In 2010, the Institute of Medicine issued a report recommending that new national sodium standards be implemented by the federal government.

ARS is USDA's chief intramural scientific research agency.

Researchers at the ARS Beltsville Human Nutrition Research Center in Beltsville, Md., developed the Automated Multiple Pass Method (AMPM), an innovative surveying tool used to obtain information on the amounts of foods and beverages consumed by a [survey](#) volunteer during a 24-hour period.

Several major food manufacturers have long been implementing sodium-reduction strategies through self-regulation. The usefulness of proposed regulatory steps will depend on accurate and practical methods to monitor the U.S. population's sodium intake.

The accuracy of sodium intake was calculated as the ratio of reported [dietary intake](#) to that estimated from [urinary sodium excretion](#). The results showed that the dietary sodium intake estimates reported by volunteers derived using the ARS survey interview method were accurate when cross-checked against the urinary sodium excretion data.

Results from this study are significant because they demonstrate that the dietary survey method is a valid tool for estimating sodium intake as well as [energy intake](#). The current ARS national food and nutrient intake survey uses the automated tool for both in-person and telephone interviews.

The automated tool will continue to accurately estimate the population's sodium intakes from foods as food composition databases produced by ARS are routinely updated to reflect changes in the salt content of foods

consumed, according to the study's lead author Donna Rhodes, a nutritionist at the Beltsville center.

The study was published in the *American Journal of Clinical Nutrition*.

Provided by Agricultural Research Service

Citation: Assessing the US population's sodium intake (2013, December 6) retrieved 25 April 2024 from <https://medicalxpress.com/news/2013-12-population-sodium-intake.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.