

# Brain structure shows who is most sensitive to pain

January 14 2014

---

Everybody feels pain differently, and brain structure may hold the clue to these differences.

In a study published in the current online issue of the journal *Pain*, scientists at Wake Forest Baptist Medical Center have shown that the brain's structure is related to how intensely people perceive pain.

"We found that individual differences in the amount of grey matter in certain regions of the brain are related to how sensitive different people are to pain," said Robert Coghill, Ph.D., professor of neurobiology and anatomy at Wake Forest Baptist and senior author of the study.

The brain is made up of both grey and [white matter](#). Grey matter processes information much like a computer, while white matter coordinates communications between the different regions of the brain.

The research team investigated the relationship between the amount of grey matter and individual differences in pain sensitivity in 116 healthy volunteers. *Pain* sensitivity was tested by having participants rate the intensity of their pain when a small spot of skin on their arm or leg was heated to 120 degrees Fahrenheit. After [pain sensitivity](#) testing, participants underwent MRI scans that recorded images of their brain structure.

"Subjects with higher pain intensity ratings had less [grey matter](#) in brain regions that contribute to internal thoughts and control of attention," said

Nichole Emerson, B.S., a graduate student in the Coghill lab and first author of the study. These regions include the [posterior cingulate cortex](#), precuneus and areas of the posterior parietal cortex, she said.

The posterior cingulate cortex and precuneus are part of the [default mode network](#), a set of connected brain regions that are associated with the free-flowing thoughts that people have while they are daydreaming.

"Default mode activity may compete with brain activity that generates an experience of pain, such that individuals with high default mode activity would have reduced sensitivity to pain," Coghill said.

Areas of the [posterior parietal cortex](#) play an important role in attention. Individuals who can best keep their attention focused may also be best at keeping pain under control, Coghill said.

"These kinds of structural differences can provide a foundation for the development of better tools for the diagnosis, classification, treatment and even prevention of [pain](#)," he said.

Provided by Wake Forest University Baptist Medical Center

Citation: Brain structure shows who is most sensitive to pain (2014, January 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2014-01-brain-sensitive-pain.html>

|   |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|