

Cancer Statistics 2014: Death rates continue to drop

January 7 2014

The annual cancer statistics report from the American Cancer Society finds steady declines in cancer death rates for the past two decades add up to a 20 percent drop in the overall risk of dying from cancer over that time period. The report, *Cancer Statistics 2014*, finds progress has been most rapid for middle-aged black men, among whom death rates have declined by approximately 50 percent. Despite this substantial progress, black men continue to have the highest cancer incidence and death rates among all ethnicities in the U.S.—about double those of Asian Americans, who have the lowest rates.

Each year, the American Cancer Society estimates the numbers of new [cancer](#) cases and deaths expected in the United States in the current year and compiles the most recent data on cancer incidence, mortality, and survival based on incidence data from the National Cancer Institute and the Centers for Disease Control and Prevention, and mortality data from the National Center for Health Statistics. The data are disseminated in two reports, *Cancer Statistics*, published in *CA: A Cancer Journal for Clinicians*, and its companion article, *Cancer Facts & Figures*.

This year's report estimates there will be 1,665,540 new cancer cases and 585,720 cancer deaths in the United States in 2014. Among men, prostate, lung, and colon cancer will account for about half of all newly diagnosed cancers, with prostate cancer alone accounting for about one in four cases. Among women, the three most common cancers in 2014 will be breast, lung, and colon, which together will account for half of all cases. Breast cancer alone is expected to account for 29% of all new

cancers among women.

The estimated 585,720 deaths from cancer in 2014 correspond to about 1,600 deaths per day. Lung, colon, prostate, and breast cancers continue to be the most common causes of [cancer death](#), accounting for almost half of the total cancer deaths among men and women. Just over one in four cancer deaths is due to lung cancer.

During the most recent five years for which there are data (2006-2010), cancer incidence rates declined slightly in men (by 0.6% per year) and were stable in women, while cancer [death rates](#) decreased by 1.8% per year in men and by 1.4% per year in women. The combined cancer death rate has been continuously declining for two decades, from a peak of 215.1 per 100,000 in 1991 to 171.8 per 100,000 in 2010. This 20 percent decline translates to the avoidance of approximately 1,340,400 cancer deaths (952,700 among men and 387,700 among women) during this time period.

The magnitude of the decline in cancer death rates from 1991 to 2010 varies substantially by age, race, and sex, ranging from no decline among white women aged 80 years and older to a 55% decline among [black men](#) aged 40 years to 49 years. Notably, black men experienced the largest drop within every 10-year age group.

"The progress we are seeing is good, even remarkable, but we can and must do even better," said John R. Seffrin, PhD, chief executive officer of the American Cancer Society. "The halving of the risk of cancer death among middle aged black men in just two decades is extraordinary, but it is immediately tempered by the knowledge that death rates are still higher among black men than white men for nearly every major cancer and for all cancers combined."

Provided by American Cancer Society

Citation: Cancer Statistics 2014: Death rates continue to drop (2014, January 7) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-01-cancer-statistics-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.