

Lowering WBC criteria lowers negative appendectomy rate

January 9 2014



(HealthDay)—Reducing the threshold of leukocytosis as a criterion for appendicitis to 8,000 to 9,000 white blood cells (WBCs) per μL reduces the rates of negative appendectomy, according to a study published online Dec. 30 in *Pediatrics*.

Maria F. Bates, M.D., from Brown University in Providence, R.I., and colleagues retrospectively reviewed data for all appendectomies performed for suspected [appendicitis](#) at a tertiary children's hospital during a 42-month period for preoperative clinical, laboratory, and radiographic variables. Sensitivities and false positive rates were determined using cutoff points on receiver operating characteristic curves for continuous variables.

The researchers found that 22 of the 847 appendectomies performed

(2.6 percent) had a pathologically normal appendix. WBC count and neutrophil count were found to be normal in 89 and 79 percent, respectively, of negative [appendectomies](#). Using WBC cutoffs of 9,000 and 8,000 per μL yielded sensitivities of 92 percent and 95 percent, respectively in receiver operating characteristic curves, and reduced the [negative appendectomy](#) rate by 77 percent and 36 percent, respectively. These expected sensitivities and specificities were confirmed in cases treated in the subsequent 12 months.

"Absence of an elevated WBC count is a risk factor for negative appendectomy," the authors write. "Withholding appendectomy for WBC counts

Citation: Lowering WBC criteria lowers negative appendectomy rate (2014, January 9) retrieved 24 April 2024 from <https://medicalxpress.com/news/2014-01-lowering-wbc-criteria-lowers-negative.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.