

Middle-school girls continue to play soccer with concussion symptoms, study shows

January 20 2014

Concussions are common among middle-school girls who play soccer, and most continue to play with symptoms, according to a study by John W. O' Kane, M.D., of the University of Washington Sports Medicine Clinic, Seattle, and colleagues.

Sports-related concussions account for 1.6 to 3.8 million injuries in the United States annually, including about 50,000 soccer-related concussions among high school [players](#). Injury-tracking systems for younger players are lacking so they are largely unstudied, according to the study background.

Using an email survey and interviews, the authors evaluated the frequency and duration of concussions in young female soccer players, as well as whether the injuries resulted in stopping play and seeking medical attention. Their study included 351 [soccer players](#) (ages 11 to 14 years) from soccer clubs in the Puget Sound region of Washington.

Among 351 players, there were 59 concussions with 43,742 athletic exposure hours. Concussion [symptoms](#) can include memory loss, dizziness, drowsiness, headache and nausea. Cumulative concussion incidence was 13 percent per season with an incidence of 1.2 per 1,000 athletic exposure hours. Symptoms lasted a median four days (average 9.4 days). Heading the ball accounted for 30.5 percent of concussions. Most players (58.6 percent) continued to play with symptoms, with almost half (44.1 percent) seeking [medical attention](#), according to the results.

The authors note that the rate of 1.3 concussions per 1,000 athletic exposure hours was higher than what has been reported in other studies of girls soccer at the high school and college levels.

"Future studies are needed to develop education strategies to ensure players understand and report [concussion](#) symptoms and that parents and coaches ensure appropriate medical evaluation and clearance before returning to play," the authors conclude. "Future studies should also compare short- and long-term outcomes for those who seek [medical care](#) and return to play according to recommended guidelines vs. those who do not seek medical care and/or return to play prematurely."

More information: *JAMA Pediatr.* Published online January 20, 2014.
[DOI: 10.1001/jamapediatrics.2013.4518](https://doi.org/10.1001/jamapediatrics.2013.4518)

Provided by The JAMA Network Journals

Citation: Middle-school girls continue to play soccer with concussion symptoms, study shows (2014, January 20) retrieved 24 April 2024 from <https://medicalxpress.com/news/2014-01-middle-school-girls-soccer-concussion-symptoms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--