

Osteoporosis / Osteopenia prevalent in chronic pancreatitis

January 31 2014



(HealthDay)—Sixty-five percent of patients with chronic pancreatitis have osteoporosis or osteopenia, according to a meta-analysis published in the February issue of *Clinical Gastroenterology and Hepatology*.

Sinead N. Duggan, Ph.D., of Trinity College Dublin, and colleagues performed a systematic review of the literature and meta-analysis to assess the prevalence of [osteoporosis](#) and osteopenia in patients with [chronic pancreatitis](#).

The researchers found that 10 studies, involving a total of 513 patients, were eligible for inclusion. The pooled prevalence rate in patients with chronic pancreatitis was 23.4 percent (95 percent confidence interval [CI], 16.6 to 32.0) for osteoporosis and 39.8 percent for osteopenia (95 percent CI, 29.1 to 51.6). Among patients with chronic pancreatitis, the

pooled prevalence rate for either osteoporosis or osteopenia was 65 percent (95 percent CI, 54.7 to 74.0).

"We recommend that bone-health assessment should form an integral part of the medical/nutritional work-up of [patients](#) with chronic pancreatitis to prevent and treat osteoporosis and, ultimately, to prevent fracture and its associated morbidity," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Osteoporosis / Osteopenia prevalent in chronic pancreatitis (2014, January 31) retrieved 11 May 2024 from <https://medicalxpress.com/news/2014-01-osteoporosis-osteopenia-prevalent-chronic-pancreatitis.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|