

# Blood clot risk remains higher than normal for at least 12 weeks after women deliver babies

February 13 2014

---

Women's blood clot risk remains elevated for at least 12 weeks after delivering a baby—twice as long as previously recognized, according to a large study presented at the American Stroke Association's International Stroke Conference 2014.

The chance of a blood clot rises during pregnancy, when platelets and other blood-clotting factors increase. The risk peaks around the time of delivery, but researchers found that afterwards it remained:

- 10.8 times higher during weeks 0-6;
- 2.2 times higher during weeks 7-12; and
- 1.4 times higher (a non-significant rise) during weeks 13-18.

By weeks 19-24, the chance of a blood clot returned to what it would have been if a woman had not delivered a baby.

Fewer than one in 10,000 women suffer a pregnancy-related blood clot six to 12 weeks after delivery.

"While rare, [blood clots](#) are a serious cause of disability and death in pregnant and post-partum women, and many members of our research team have cared for young women with these complications," said Hooman Kamel, M.D., lead researcher and assistant professor in the Department of Neurology and the Brain and Mind Research Institute of

Weill Cornell Medical College in New York City.

"If you have recently delivered a baby, seek medical attention if you develop symptoms such as: chest pain or pressure; difficulty breathing; swelling or pain in one leg; sudden severe headache; or sudden loss of speech, vision, balance, or strength on one side of your body."

The researchers calculated the odds using data on 1,687,930 women admitted for labor and delivery at a California hospital in 2005-10. Of those, 1,015 women had a clot during the following 1.5 years. These included [women](#) with clots that interrupted blood flow in the brain (stroke), heart (heart attack), lungs (pulmonary embolism) or limbs ([deep vein thrombosis](#)).

"Clinicians should consider our results when caring for high-risk postpartum patients, such as those with previous clots, or postpartum patients with symptoms concerning for thrombosis," Kamel said.

Provided by American Heart Association

Citation: Blood clot risk remains higher than normal for at least 12 weeks after women deliver babies (2014, February 13) retrieved 6 May 2024 from <https://medicalxpress.com/news/2014-02-blood-clot-higher-weeks-women.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--