


Intensive lifestyle reverses glucose abnormalities in youth

February 1 2014



(HealthDay)—An intensive lifestyle intervention can reverse early glucose abnormalities in obese youth, according to a study published in the February issue of *Diabetes Care*.

Mary Savoye, R.D., C.D.E., from Yale University in New Haven, Conn., and colleagues randomized obese adolescents (10 to 16 years old; Tanner stage >2) with elevated [oral glucose tolerance test](#) (OGTT) two-hour blood glucose (130 to 199 mg/dL) to either the Bright Bodies (BB) Healthy Lifestyle Program, including exercise and nutrition/behavior modification, or standard clinical care (CC). At baseline and at six months, OGTTs as well as cardiovascular and anthropometric assessments were conducted.

The researchers found that BB participants had greater reductions in two-

hour glucose compared to CC participants ($P = 0.005$). Furthermore, there was greater conversion to

Citation: Intensive lifestyle reverses glucose abnormalities in youth (2014, February 1) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2014-02-intensive-lifestyle-reverses-glucose-abnormalities.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.