

## Rare bacteria outbreak in cancer clinic tied to lapse in infection control procedure

February 12 2014

---

Improper handling of intravenous saline at a West Virginia outpatient oncology clinic was linked with the first reported outbreak of *Tsukamurella* spp., gram-positive bacteria that rarely cause disease in humans, in a new report from the Centers for Disease Control and Prevention (CDC). The report was published in the March issue of *Infection Control and Hospital Epidemiology*, the journal of the Society for Healthcare Epidemiology of America.

"This outbreak illustrates the need for outpatient clinics to follow proper [infection control guidelines](#) and medication preparation practices to minimize the risk of infection for patients with weakened immune systems," said Isaac See, MD, lead author of the study. "A combination of careful descriptive epidemiology with attention to outlier cases, direct observations, and analytic studies were needed to support this investigation, which pointed to deficiencies in medication preparation practices as the cause of these unusual infections."

From September 2011-May 2012, 15 immunocompromised patients developed *Tsukamurella* bloodstream infections. All patients had received a diagnosis of malignancy, and had an indwelling central line, although central line types varied. A case-control study determined that the only risk factor for developing *Tsukamurella* infection was the receipt of saline flush, prepared by the clinic staff from large preservative-free bags of saline, from the clinic during September-October 2011.

Investigations by the West Virginia Bureau of Public Health (WVBPH) and the CDC found several lapses in infection control procedures relating to the care of long-term intravenous catheters and preparation of chemotherapy for patients at the clinic. These investigations also suggested that saline flush syringes were the likely source of infection.

Following the recommendations of WVBPH and CDC, the clinic instituted several changes to its infection prevention and control practices; including using pre-packaged manufactured saline flushes. After the clinic changed this practice, *Tsukamurella* bloodstream infections stopped occurring, further supporting the saline flush as the source of infection.

To help outpatient oncology facilities establish appropriate infection control strategies, the CDC developed a basic [infection control](#) plan tailored to these settings outlining key policies and procedures needed to meet minimal requirements for patient safety. These include the proper use and handling of injectable medications and correct procedures for assessing central lines. Outpatient oncology facilities without an existing plan are encouraged to use this document as a starting point.

**More information:** Isaac See, Duc B. Nguyen, Somu Chatterjee, Thein Shwe, Melissa Scott, Sherif Ibrahim, Heather Moulton-Meissner, Steven McNulty, Judith Noble-Wang, Cindy Price, Kim Schramm, Danae Bixler, Alice Y. Guh. "Outbreak of *Tsukamurella* spp. Bloodstream Infections among Patients of an Oncology Clinic—West Virginia, 2011-2012." *Infection Control and Hospital Epidemiology* 35:3 (March 2014).

Provided by Society for Healthcare Epidemiology of America

Citation: Rare bacteria outbreak in cancer clinic tied to lapse in infection control procedure (2014, February 12) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-02-rare-bacteria-outbreak-cancer-clinic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.