

Using susceptibility-weighted imaging to study concussion in college ice hockey players

February 4 2014

Using susceptibility-weighted imaging (SWI), researchers have identified microstructural changes in the brains of male and female college-level ice hockey players that could be due to concussive or subconcussive trauma. Until now, SWI has been used to detect signs of more severe cases of traumatic brain injury (TBI). This is the first time SWI has been used to detect signs of concussion (or mild TBI), and the first time it has been used to detect changes in the brain prospectively over an entire sports season in athletes of both sexes. The researchers, hailing from several institutions,¹ describe and discuss their findings in "Hockey Concussion Education Project, Part 1. Susceptibility-weighted imaging study in male and female ice hockey players over a single season. Clinical article," by Karl G. Helmer and colleagues, published online today, ahead of print, in the *Journal of Neurosurgery*. Two additional papers from the Hockey Concussion Education Project (HCEP) on imaging techniques used to detect and follow concussion's effects are also being published today.²

Susceptibility-weighted imaging (SWI) is a high-resolution three-dimensional gradient-echo magnetic resonance imaging (MRI) technique that is particularly sensitive to the presence of blood and blood-breakdown products. SWI can be used to detect cerebral microbleeds, which are small hemorrhages that appear hypointense (contrasting to the appearance of other parts of the brain) on T2*-weighted magnetic resonance images. These microbleeds are often the result of traumatic brain injuries (TBIs). Until now SWI has been used to detect microbleeds larger than 5 millimeters—often a sign of severe TBI. In

this study, Helmer and colleagues used SWI to detect clusters of much smaller cerebral microbleeds—ones that are too small to be seen with the naked eye but could represent acute and chronic injuries to the brain due to concussion or subconcussive events.

The authors report findings in college-level ice hockey players (25 men and 20 women) who initially agreed to participate in the study. These results document the innovative SWI analysis of the participating athletes prospectively during the hockey season, including pre- and post-season testing. Some of these athletes reported a history of concussion prior to the study period and some reported receiving no such injuries. During the season, 5 male and 6 [female athletes](#) sustained concussions that were identified and diagnosed by independent (study) medical specialists that attended each game. These athletes underwent additional SWI at 72 hours, 2 weeks, and 2 months after their injuries as well as serial clinical and neuropsychological evaluations. The researchers identified small clusters of hypointensity, which can be related to concussion (mild TBI), on SWIs obtained at time points in all participants. To compare the extent of these hypointense regions within and between individual athletes and groups of athletes, the researchers calculated a hypointensity burden (HIB) metric for each athlete at each SWI time point.

Helmer and colleagues report no significant change in the HIB value between the beginning and end of the hockey season in either male or female athletes who did not sustain a concussion. When the researchers compared HIB values in nonconcussed athletes according to sex, however, they did identify significant differences both at the beginning and end of the season, with male athletes having significantly higher mean HIB values than female athletes at both time points.

Among the athletes who did sustain a concussion during the season, the researchers report a statistically significant difference between the HIB

value determined at the beginning of the season and that determined 2 weeks after concussive injury in male athletes and a similar—albeit not significant—difference in the HIB value during the same period in female athletes. These changes in HIB values point toward damage sustained at the time of impact as well and may indicate later effects of concussion. It is not unusual for a TBI to produce a two-pronged effect: immediate damage from the impact and secondary damage from injury-induced metabolic changes in the brain. In concussed athletes, the researchers also identified a difference in HIB values between the sexes, with male athletes having higher mean HIB values than female athletes at all SWI time points throughout the season.

The authors note that not all clusters of hypointensity seen on susceptibility-weighted images may represent small microbleeds; some may prove to be false-positive findings. However, given changes in the HIB metric following concussion, the researchers believe that SWI may be a useful diagnostic tool in detecting the effects of concussive impacts over time.

As to the differences in the HIB metric between the sexes, the researchers hypothesize various reasons why this occurred, including the possibility that the male athletes may have accumulated more extensive injuries from pre-study concussive and subconcussive impacts. Additional studies in a greater number of participants are called for, because these may lead to a clearer description of why the HIB metric differed between the sexes in this study.

In summarizing the value of SWI for detecting injuries due to [concussion](#) in [athletes](#) participating in contact sports, the researchers state: "Data acquired using this method could be used for the monitoring of players throughout their careers and could lead to improved diagnoses and return-to-play guidelines."

When asked to comment on the importance of this study, Dr. Paul Echlin, senior co-author and primary HCEP investigator in this study, stated, "This study contributes to converging objective evidence concerning the acute and chronic effects of repetitive brain injury in the sports. Although future studies are warranted to further validate these initial findings (as well those found in the two accompanying HCEP studies), a cultural shift must be considered toward both the permitted violence that underlies many of the sport-related brain injuries and the high incidence of head impacts that occur in the games which our children play."

More information: Helmer KG, Pasternak O, Fredman E, Preciado RI, Koerte IK, Sasaki T, Mayinger M, Johnson AM, Holmes JD, Forwell LA, Skopelja EN, Shenton ME, Echlin PS: Hockey Concussion Education Project, Part 1. Susceptibility-weighted imaging study in male and female ice hockey players over a single season. Clinical article. Journal of Neurosurgery, published online, ahead of print, February 4, 2014; DOI: 103171/2013.12.JNS132093

Provided by Journal of Neurosurgery

Citation: Using susceptibility-weighted imaging to study concussion in college ice hockey players (2014, February 4) retrieved 23 April 2024 from <https://medicalxpress.com/news/2014-02-susceptibility-weighted-imaging-concussion-college-ice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.