

Well-child visits linked to more than 700,000 subsequent flu-like illnesses

February 12 2014

New research shows that well-child doctor appointments for annual exams and vaccinations are associated with an increased risk of flu-like illnesses in children and family members within two weeks of the visit. This risk translates to more than 700,000 potentially avoidable illnesses each year, costing more than \$490 million annually. The study was published in the March issue of *Infection Control and Hospital Epidemiology*, the journal of the Society for Healthcare Epidemiology of America.

"Well child visits are critically important. However, our results demonstrate that healthcare professionals should devote more attention to reducing the risk of spreading infections in waiting rooms and clinics. Infection control guidelines currently exist. To increase patient safety in outpatient settings, more attention should be paid to these guidelines by healthcare professionals, patients, and their families," said Phil Polgreen, MD, MPH, lead author of the study.

Researchers from the University of Iowa used data from the Agency for Healthcare Research and Quality's (AHRQ) Medical Expenditure Panel Survey to examine the healthcare trends of 84,595 families collected from 1996-2008. Included in the analysis were demographic, office-based, emergency room, and outpatient cases records. After controlling for factors, such as the presence of other children, insurance, and demographics, the authors found that well-child visits for children younger than six years old increased the probability of a flu-like illness in these children or their families during the subsequent two weeks by



3.2 percentage points.

This incremental risk could amount to more than 700,000 avoidable cases of flu-like illness each year and \$492 million in direct and indirect costs, based on established estimates for outpatient influenza.

In a commentary accompanying the study, Lisa Saiman, MD, notes, "The true cost of flu-like illnesses are much higher since only a fraction result in ambulatory visits and many more cases are likely to result in missed work or school days. Furthermore, these flu-like illness visits are associated with inappropriate antimicrobial use."

The authors stress the importance of infection prevention and control in ambulatory settings, suggesting pediatric clinics follow recommended guidelines that include improving environmental cleaning, cough etiquette, and hand hygiene compliance.

"Even with interventions, such as the restricted use of communal toys or separate sick and well-child waiting areas, if hand-hygiene compliance is poor, and potentially infectious patients are not wearing masks, preventable infections will continue to occur," said Polgreen.

Provided by Society for Healthcare Epidemiology of America

Citation: Well-child visits linked to more than 700,000 subsequent flu-like illnesses (2014, February 12) retrieved 20 April 2024 from https://medicalxpress.com/news/2014-02-well-child-linked-subsequent-flu-like-illnesses.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.