

Young steroid users at increased risk of heart disease

February 27 2014, by Marion Downey



Steroid abuse is associated with increased risk of heart disease in otherwise healthy young men, an Australian study of deaths involving the drugs has found.

Researchers at the National Drug and Alcohol Research Centre (NDARC) at UNSW and the New South Wales Department of Forensic Medicine examined all 24 steroid-related deaths in NSW between 1996 - 2012. Extensive heart disease, including left ventricular hypertrophy and thickening of the arteries, was present in half the cases. This is notable

considering the sample comprised exclusively of men aged 22-48 years, with an average age of 32.

Although all the cases showed evidence of steroid use, these were not the direct cause of death. All but one of the sample were polydrug users and accidental drug toxicity accounted for two thirds of the deaths, either alone or in combination with [heart disease](#). Violent deaths – either suicide or homicide – accounted for a quarter of cases.

Lead researcher Professor Shane Darke of NDARC said the findings added to emerging evidence that steroid users are often polydrug users, and that steroid and psychostimulant use may damage the heart.

"Nearly all of the 24 deceased men that we examined in this study showed the classic signs of [steroid abuse](#), such as overdeveloped muscles and testicular atrophy," Darke said.

"We also found that nearly all of the individuals who had been using [steroids](#) together with psychostimulants had a lot of damage to their cardiovascular system. It appears that combined stimulant and steroid use increases a person's risk of developing [cardiovascular disease](#).

"Ironically many steroid users are also keen health and fitness enthusiasts and this is borne out by our sample, which included personal trainers, body builders and security guards.

"But the vast majority of users are unaware of the health dangers of their lifestyle.

"Cardiovascular disease, particularly in such young people, signifies the very opposite of good health," Darke said.

He added that many young male users are in danger of reproductive

health issues attributable to testicular atrophy.

Among the psychoactive drugs detected were psychostimulants (present in 66.7% of cases), followed by benzodiazapines (45.8%), opioids (37.5%) and alcohol (25%). Psychostimulant toxicity was the direct cause of death in eight of the 24 deaths and opioid overdose was the direct cause in seven. Over half of the cases (54%) showed evidence of recent injecting drug use.

Toxicology results showed testosterone was the most commonly used steroid among the 24 men. Two thirds of the men were employed at their time of death.

More information: *Journal of Forensic Science*
newsroom.unsw.edu.au/sites/default/files/documents/Paper%20published%20online%2019%20Feb.pdf

Provided by University of New South Wales

Citation: Young steroid users at increased risk of heart disease (2014, February 27) retrieved 24 April 2024 from
<https://medicalxpress.com/news/2014-02-young-steroid-users-heart-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--