


New evidence links smoking to postmenopausal breast cancer risk

March 19 2014



Postmenopausal women who smoke or have smoked in the past may have an increased risk of breast cancer compared with women who have never smoked according to a new study published in the *British Journal of Cancer*.

Researchers from the U.S. National Cancer Institute in Bethesda, Maryland found that [postmenopausal women](#) who reported they smoke were 19 per cent more likely to develop [breast cancer](#) compared with

[women](#) who never smoked.

And those who said they were former smokers were seven per cent more likely to be diagnosed with the disease than women who had never smoked.

These results held true even after accounting for alcohol consumption, which is known to increase the risk of breast cancer, and is more common among smokers.

The researchers asked around 186,000 American women aged between 50 and 71 in the mid-1990s about their [smoking](#) and alcohol use, and their medical and reproductive history. They then tracked these women for around 10 years. Of these around 7,500 developed breast cancer in that time.

Sarah Nyante, one of the study authors based at the U.S. National Cancer Institute in Bethesda, Maryland, said: "Our study adds to a growing body of evidence that suggests an association between cigarette smoking and increased breast cancer risk.

"Previous studies have investigated this relationship, but questions remained regarding the extent to which other [breast cancer risk](#) factors, such as alcohol intake, might influence the results. More work is now needed to understand the mechanisms behind the link between smoking and breast cancer in postmenopausal women."

Lifestyle factors such as being overweight, drinking alcohol and not keeping physically active can all increase the risk of breast cancer among postmenopausal women – but the role of smoking is more difficult to unravel.

We do know that smoking is the biggest preventable causes of cancer

worldwide. It is estimated that one in two regular cigarette smokers will eventually be killed by their tobacco addiction. Over the last 50 years, it is estimated that 6.5 million people in the UK have died from tobacco-related diseases.

Dr Julie Sharp, Cancer Research UK's head of health information, said: "Evidence remains inconsistent as to whether smoking causes breast cancer after as well as before the menopause, but this study suggests it may increase a postmenopausal woman's risk of breast cancer if she smokes or has smoked in the past.

"But there is strong evidence that smoking increases your risk for many other types of cancer including lung, larynx, oesophageal, oral cavity and pharynx, bladder, pancreas, kidney, liver, stomach and bowel. The best thing you can do if you are a smoker is give up. Quitting is not easy but, given that smokers lose an average of 10 years of life compared to non-smokers, the benefits are huge."

More information: "Cigarette smoking and postmenopausal breast cancer risk in a prospective cohort." S J Nyante, G L Gierach, C M Dallal, N D Freedman, Y Park, K N Danforth, A R Hollenbeck and L A Brinton *British Journal of Cancer* , 18 March 2014 [DOI: 10.1038/bjc.2014.132](https://doi.org/10.1038/bjc.2014.132)

Provided by Cancer Research UK

Citation: New evidence links smoking to postmenopausal breast cancer risk (2014, March 19) retrieved 10 April 2024 from <https://medicalxpress.com/news/2014-03-evidence-links-postmenopausal-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.