

Iron deficiency important to assess in children adopted from institutional settings

March 6 2014

A new longitudinal study finds that children who spent more time in institutional settings (like orphanages) prior to adoption, and had more severe iron deficiency at the time of adoption, were more likely to have lower IQs and poorer higher-order thinking skills a year later. The study—which followed children adopted into US families from Africa, Asia, Eastern Europe and Asia—suggests that iron supplements and cognitive interventions could be helpful in counteracting these effects.

Iron deficiency predicts lower IQ scores and poor higher-order [thinking skills](#) in [children](#) adopted from institutional settings like orphanages, according to a new longitudinal study.

The study analyzed data on 55 children adopted from international institutions, with a focus on nutritional status. Conducted by researchers at the University of Minnesota, the research appears in the journal *Child Development*.

Children with more severe [iron deficiency](#) when they were adopted and who had spent more time in an institution before they were adopted were more likely to have lower IQ and poorer higher-order thinking skills (such as the ability to plan and solve problems), the study found. Iron deficiency affected higher-order thinking skills independent of its effect on general cognitive ability, as measured by the IQ test. Even less severe levels of iron deficiency were associated with poorer [cognitive functioning](#) (examples of cognitive functioning include planning behavior, storing information in memory and then acting on that

information, and thinking flexibly about situations).

"Our study shows that both the duration of institutional care and nutrition significantly affect children's cognitive functioning even after they are placed in a nurturing home," according to Jenalee Doom, graduate student, and Megan Gunnar, professor of child psychology, at the University of Minnesota, two of the study's authors.

"These findings are important for pediatricians, psychologists, and other specialists working with children adopted internationally because they highlight the importance of examining both nutritional deficiencies and children's pre-adoptive experience during evaluation and treatment."

Specifically, they note, children can receive iron supplements or [cognitive interventions](#) that address each child's specific needs. Common interventions for internationally adopted children include those that address sensory integration disorder and attention deficit hyperactivity disorder. Parents should be informed about adoption nutrition so they make sound nutritional choices to optimize their children's development. Organizations that specialize in nutrition in orphanages will benefit from an understanding of which nutrients are vital for cognitive functioning and what types of nutrient deficiencies are commonly found in children adopted internationally who had previously lived in institutional settings.

Even children who aren't adopted may benefit from careful monitoring of iron status in early childhood, given that deficiency significantly predicted cognitive outcomes independent of how long the children in this study had been institutionalized, the authors note.

The study examined IQ, higher-order thinking skills, and iron deficiency in children adopted into U.S. families from Africa, Asia, Eastern Europe, and Latin America. Previous work has looked at the effect of early deprivation on cognitive development among children living in

institutions such as orphanages or hospitals. But few studies have examined whether these children suffer from nutritional deficiencies that would help explain disruptions in cognitive functioning.

Researchers examined medical records of 55 children (then ages 17 to 36 months) soon after they were adopted and classified them as having normal iron levels, less severe iron deficiency, or severe iron deficiency, characterized as anemia. About a year later, when the children were 2 to 4 years old, they were given an IQ test and tests that measure higher-order thinking skills. These skills, also called executive function, include the ability to delay a reward, store and manage information to accomplish a task, and think flexibly about a task.

Provided by Society for Research in Child Development

Citation: Iron deficiency important to assess in children adopted from institutional settings (2014, March 6) retrieved 23 April 2024 from <https://medicalxpress.com/news/2014-03-iron-deficiency-important-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.