

# Pre-op pregabalin best for pain in spinal surgery

March 28 2014

---



(HealthDay)—Preoperative pregabalin is superior to either gabapentin or placebo for the relief of pain in patients undergoing lumbar discectomy, according to a study published in the March 15 issue of *Spine*.

Gurjeet Khurana, M.D., from HIHT University in Dehradun, India, and colleagues compared 300 mg of [gabapentin](#) (Group A), 75 mg of pregabalin (Group B), or placebo (Group C) in 90 patients (aged 18 to 75 years) undergoing spinal surgery. One dose was given one hour before surgery and at eight hour intervals for seven days afterwards. Postoperative static and dynamic pain and functional outcome were assessed using a visual analogue scale, the Prolo functional and economic score, and the Oswestry Disability Index score, both preoperatively and three months postoperatively.

The researchers found that subjects receiving pregabalin showed consistently reduced static and dynamic pain intensity and also required a lesser amount of rescue drug throughout the [postoperative period](#). At all time intervals there was a statistically significant difference (P

"Preoperative pregabalin administration is associated with less pain intensity and improved functional outcomes three months after lumbar discectomy followed by gabapentin and then [placebo](#)," the authors write.

**More information:** [Full Text](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Pre-op pregabalin best for pain in spinal surgery (2014, March 28) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-03-pre-op-pregabalin-pain-spinal-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--