

New approach for surgery patients cuts hospital stays and costs

April 24 2014

Changes in managing patients before, during and after colorectal surgery cut hospital stays by two days and reduced readmission rates, according to researchers who led a study of the approach at Duke University Hospital.

The practice, called enhanced recovery, is easier on [patients](#) before surgery, doing away with the fasting period and bowel evacuation that are typically prescribed. After surgery, patients are encouraged to eat and move about as soon as possible, leading to faster recoveries.

Among findings published in the May 2014 issue of the journal *Anesthesia & Analgesia*, the researchers reported that the enhanced recovery approach used for colorectal surgery cut hospital admissions from an average seven days to five, and reduced the rate of readmissions by half.

"Enhanced recovery is about change management," said senior author Tong J. Gan, M.D., MHS, professor of anesthesiology at Duke. "It's getting the team together, including nurses, anesthesiologists, surgeons and patients, with everyone understanding the expectations of how to do things differently and improve patient care."

Gan and colleagues collected data from 241 consecutive patients at Duke University Hospital undergoing open or laparoscopic colorectal surgery during two time periods: before the enhanced recovery approach was implemented, and after. Ninety-nine patients were studied in the

traditional approach, and 142 using enhanced recovery.

With traditional perioperative care – defined as the care provided through a patient's hospitalization for surgery – few procedures are standardized, but patients are typically told to fast the night before and undergo laxative treatments, and then are not given food or drink after surgery until bowel sounds are restored, sometimes several days later. Additionally, traditional perioperative care includes a variety of different anesthesia regimens, fluid management and pain control, depending on the surgical team's preferences.

Under traditional perioperative care, patients who experience pain, stress, immobilization, and postoperative constipation can remain in the hospital for 10 days or more, with complication rates of up to 48 percent. Such complications can be expensive, estimated at an average of \$10,000.

Enhanced recovery after surgery (ERAS), which has gained favor in Europe but has not been widely used in the United States, aims to standardize perioperative care using procedures backed by scientific evidence that demonstrate their benefits.

In the Duke study, patients in the ERAS group were educated about what they should expect. Routine bowel preparation was not performed, and patients were allowed to drink clear fluids until 3 hours before their surgeries, notably a sports drink.

All of the ERAS patients received an epidural as well as non-opioid painkillers to reduce opioid side effects such as nausea, vomiting, constipation, urinary retention and drowsiness. They then underwent general anesthesia. After surgery, the patients transitioned to oral acetaminophen or other non-steroidal anti-inflammatory drugs, plus oral opioids, if necessary, after about 72 hours. Patients were also

encouraged to drink liquids and get out of bed on the day of surgery, and for at least six hours every subsequent day.

"We have shown that providing care within an ERAS pathway, we are increasing the quality of care for patients while at the same time reducing complications and medical costs," said co-author Timothy Miller, assistant professor of anesthesiology at Duke. "I believe that going forward, enhanced recovery care could become the new standard for best practice."

Gan said the researchers saved costs for about 85 percent of the ERAS patients, at about \$2,000 per patient. He said the ERAS approach could be used for numerous surgeries; it has been expanded at Duke University Hospital to include bladder, pancreas and liver surgeries.

"Our study demonstrates that the patient-focused enhanced recovery approach uses evidence-based concepts to decrease perioperative stress and improve outcomes from [surgery](#)," said lead colorectal surgeon and co-author, Julie K. Marosky Thacker, M.D.

Provided by Duke University Medical Center

Citation: New approach for surgery patients cuts hospital stays and costs (2014, April 24)
retrieved 19 April 2024 from
<https://medicalxpress.com/news/2014-04-approach-surgery-patients-hospital.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--