

Stroke risk reduced if brain blood vessel disorder is left alone

April 29 2014

Treating patients who suffer from a common condition that affects blood vessels in the brain increases their risk of stroke, a study has found.

People with a condition known as [arteriovenous malformation](#) (AVM) – which causes [blood vessels](#) in the [brain](#) to tangle – have a better outcome if doctors treat their symptoms only and not the AVM.

A team of doctors looked at the long-term outcome of patients with the condition, which is caused by abnormal connections between the arteries and veins in the brain.

They found that, over a 12 year period, patients who chose not to be treated for their condition were less likely to have a stroke or die from related causes. These patients were also less likely to suffer sustained disability compared with those who opted for an intervention to treat the tangles.

This is the first study to compare the risks and benefits of treatment for AVM in the long term. The findings build on previous research that reported an increased risk of stroke in the first three years after treatment for AVM.

AVM affects around 1 in 2000 people. Although most people with the condition can lead relatively normal lives, they live with the risk that the tangles can burst and bleed into the brain at any time, causing a stroke.

Around one in every hundred AVM patients suffers a stroke each year.

In some cases doctors can surgically remove the tangle or block the blood vessels involved to reduce the risk of bleeding. However, treatment can sometimes increase the chances that the tangle will burst and bleed into the brain, resulting in a [stroke](#).

Professor Rustam Al-Shahi Salman, MRC Senior Clinical Fellow at the University of Edinburgh and a consultant neurologist, said: "Many [patients](#) feel that living with AVM is like living with a time bomb in your head that could explode at any time. Patients and their doctors face difficult choices when deciding whether or not to pursue treatment. We have found that, for most people whose AVM has not caused a bleed, the risks of [treatment](#) exceed the risks of leaving it alone over 12 years."

The study is published in the *Journal of the American Medical Association*.

Provided by University of Edinburgh

Citation: Stroke risk reduced if brain blood vessel disorder is left alone (2014, April 29)
retrieved 26 April 2024 from

<https://medicalxpress.com/news/2014-04-brain-blood-vessel-disorder-left.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--