


Controlling brain waves to improve vision

April 24 2014



Position of each of the light sources (red) and detectors (yellow) across both sessions, as well as three-dimensional reconstruction of the estimated light path of each optical channel between a source and detector, shown as green ellipsoids for each channel, both co-registered with the individual's structural magnetic resonance image (MRI) for a representative subject. Background is a long-exposure photograph of the near-infrared optic fibers carrying light from the imaging instrument. Credit: Kyle Mathewson, Beckman Institute for Advanced Science and Technology

Have you ever accidentally missed a red light or a stop sign? Or have you heard someone mention a visible event that you passed by but totally missed seeing?

"When we have different things competing for our attention, we can

only be aware of so much of what we see," said Kyle Mathewson, Beckman Institute Postdoctoral Fellow at the University of Illinois. "For example, when you're driving, you might really be concentrating on obeying traffic signals."

But say there's an [unexpected event](#): an emergency vehicle, a pedestrian, or an animal running into the road—will you actually see the unexpected, or will you be so focused on your initial task that you don't notice?

"In the car, we may see something so brief or so faint, while we're paying attention to something else, that the event won't come into our awareness," says Mathewson. "If you present this scenario hundreds of times to someone, sometimes they will see the unexpected event, and sometimes they won't because their brain is in a different preparation state."

By using a novel technique to test [brain waves](#), Mathewson and colleagues at the Beckman Institute for Advanced Science and Technology at the University of Illinois are discovering how the brain processes external stimuli that do and don't reach our awareness. A paper about their results, "Dynamics of Alpha Control: Preparatory Suppression of Posterior Alpha Oscillations by Frontal Modulators Revealed with Combined EEG and Event-related Optical Signal," published this month in the *Journal of Cognitive Neuroscience*, reveals how [alpha waves](#), typically thought of as your brain's electrical activity while it's at rest, can actually influence what we see or don't see.

The researchers used both electroencephalography (EEG) and the event-related optical signal (EROS), developed in the Cognitive Neuroimaging Laboratory of Gabriele Gratton and Monica Fabiani, professors of psychology and members of the Beckman Institute's Cognitive Neuroscience Group, and authors of the study.

While EEG records the [electrical activity](#) along the scalp, EROS uses infrared light passed through optical fibers to measure changes in optical properties in the active areas of the cerebral cortex. Because of the hard skull between the EEG sensors and the brain, it can be difficult to find exactly WHERE signals are produced. EROS, which examines how light is scattered, can noninvasively pinpoint activity within the brain.

"EROS is based on near-infrared light," explained Fabiani and Gratton via email. "It exploits the fact that when neurons are active, they swell a little, becoming slightly more transparent to light: this allows us to determine when a particular part of the cortex is processing information, as well as where the activity occurs."

This allowed the researchers to not only measure activity in the brain, but also allowed them to map where the alpha oscillations were originating. Their discovery: the alpha waves are produced in the cuneus, located in the part of the brain that processes visual information.

The alpha can inhibit what is processed visually, making it hard for you to see something unexpected.

By focusing your attention and concentrating more fully on what you are experiencing, however, the executive function of the brain can come into play and provide "top-down" control—putting a brake on the alpha waves, thus allowing you to see things that you might have missed in a more relaxed state.

"We found that the same brain regions known to control our attention are involved in suppressing the alpha waves and improving our ability to detect hard-to-see targets," said Diane Beck, a member of the Beckman's Cognitive Neuroscience Group, and one of the study's authors.

"Knowing where the waves originate means we can target that area

specifically with electrical stimulation" said Mathewson. "Or we can also give people moment-to-moment feedback, which could be used to alert drivers that they are not paying attention and should increase their focus on the road ahead, or in other situations alert students in a classroom that they need to focus more, or athletes, or pilots and equipment operators."

The study examined 16 subjects and mapped the electrical and optical data onto individual MRI [brain](#) images.

Provided by Beckman Institute for Advanced Science and Technology

Citation: Controlling brain waves to improve vision (2014, April 24) retrieved 20 March 2024 from <https://medicalxpress.com/news/2014-04-brain-vision.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--