

Consuming a high-fat diet is associated with increased risk of certain types of breast cancer

April 9 2014

High total and saturated fat intake were associated with greater risk of estrogen receptor- and progesterone receptor-positive (ER+PR+) breast cancer (BC), and human epidermal growth factor 2 receptor-negative (HER2-) disease, according to a new study published April 9 in the *Journal of the National Cancer Institute*.

Published data from epidemiological and case-control studies on the association between high [fat intake](#) and BC risk have been conflicting, which may be attributable to difficulties obtaining accurate information on fat intake and because of limited heterogeneity of intake within a specific geographic area from which the study cohorts live. Furthermore, BC is now classified clinically into subtypes by ER, PR, and HER2 expression status and each subtype has its own prognosis and set of [risk factors](#), which may also contribute to the inconsistencies in the published reports on this relationship.

Sabina Sieri, Ph.D., from the Epidemiology and Prevention Unit of the Department of Preventive & Predictive Medicine at Fondazione IRCCS Istituto Nazionale Dei Tumori in Milan Italy, and colleagues prospectively analyzed data from 10,062 BC patients from the EPIC study with 11.5 years of follow-up. The EPIC cohort study consisted of 337,327 women living in 10 European countries, which creates a heterogeneous cohort both in terms of geography-related dietary fat intake patterns and in terms of molecular subtype. To correct the dietary

questionnaire data for measurement errors, intake data were calibrated with standardized 24-hour dietary recall interviews on administered to a random sample of 8% of the cohort. Cox proportional hazard modeling included various known risk factors as covariables.

The authors report high total and [saturated fat](#) intake were associated with greater risk of ER+PR+ BC. High saturated fat intake was also associated with greater risk of HER2- disease. The authors conclude, "a high-fat diet increases BC risk and, most conspicuously, that high saturated fat intake increases risk of receptor-positive disease, suggesting saturated fat involvement in the etiology of receptor-positive BC."

Provided by Oxford University Press

Citation: Consuming a high-fat diet is associated with increased risk of certain types of breast cancer (2014, April 9) retrieved 20 March 2024 from <https://medicalxpress.com/news/2014-04-consuming-high-fat-diet-breast-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--