


Inverse link for carotenoid intake, benign breast disease

April 8 2014



(HealthDay)—For adolescent girls, β -carotene intake is inversely associated with the risk of benign breast disease (BBD), according to a study published online April 7 in *Pediatrics*.

Caroline E. Boeke, Sc.D., from the Harvard School of Public Health in Boston, and colleagues examined adolescent carotenoid intake in relation to BBD in 6,593 [adolescent girls](#) in the prospective Growing Up Today Study cohort. Food frequency questionnaires were administered in 1996, 1997, and 1998 to assess intake of α -[carotene](#), β -carotene, β -cryptoxanthin, lutein/zeaxanthin, and lycopene. In questionnaires administered in 2005, 2007, and 2010, girls reported on biopsy-confirmed BBD.

The researchers identified an inverse correlation between β -carotene and

BBD, with a multivariate-adjusted odds ratio of 0.58 comparing the highest to the lowest quartile (P trend = 0.03). Inverse associations were also observed for intake of α -carotene and lutein/zeaxanthin with BBD, but these were not statistically significant.

"Adolescent carotenoid intake may be associated with lower BBD risk; these findings warrant further study," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Inverse link for carotenoid intake, benign breast disease (2014, April 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2014-04-inverse-link-carotenoid-intake-benign.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--