

Microfluidic technology reveals potential biomarker for early pancreatic cancer

April 29 2014

Cancer cells are on the move in the bloodstream in the very early stage of pancreatic cancer, and can be detected before cancer is diagnosed, according to research by the University of Michigan Health System.

In a study of 51 patients, researchers used a state-of-the-art microfluidic device to capture circulating pancreas [epithelial cells](#) in 33 percent of patients with early pancreatic lesions and no clinical diagnosis of cancer.

The findings, published in *Gastroenterology*, suggest that circulating [pancreas cells](#) (CPCs) seed the [bloodstream](#) before tumors can be detected using current clinical tests such as CT and MRI scans. The data suggest that the detection of pancreas cells in the blood may be an early sign of cancer.

"While there is much work that still needs to be done, there is great potential for using this technology to identify who is most at risk for developing [pancreatic cancer](#)," says lead author Andrew Rhim, M.D., an assistant professor of internal medicine at the U-M Health System and gastroenterologist at the U-M Comprehensive Cancer Center's Multidisciplinary Pancreatic Cancer Clinic.

One reason the outlook for pancreatic cancer is poor is that very few of the cancers are found early. Patients usually have no symptoms until the cancer has spread to other organs. There are no blood tests to find early cancers of the pancreas.

It is thought that [cancer cells](#) seed the bloodstream late in cancer progression, when large tumors are present. But based on studies in animal models of pancreatic cancer, U-M researchers and colleagues theorized that dissemination happens before we tumors can be detected.

They studied three groups of patients: those who were cancer-free, those diagnosed with precancerous cystic lesions, and patients with pancreatic cancer. Researchers captured circulating epithelial cells in 73 percent of patients with cancer. None of the 19 cancer-free patients without lesions had these cells in their bloodstream.

"Studies are underway to interrogate the genomic signature of circulating pancreas cells from [patients](#) with precancerous cystic lesions," says Rhim.

"If these cells represent the earliest forms of cancer, we predict they would possess many of the genetic anomalies we typically see in pancreas tumors."

More information: "Detection of Circulating Pancreas Epithelial Cells in Patients With Pancreatic Cystic Lesions," *Gastroenterology*, Vol. 146, Issue 3.

Provided by University of Michigan Health System

Citation: Microfluidic technology reveals potential biomarker for early pancreatic cancer (2014, April 29) retrieved 19 April 2024 from <https://medicalxpress.com/news/2014-04-microfluidic-technology-reveals-potential-biomarker.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.