

Proof that antidepressants and breastfeeding can mix

April 10 2014, by David Ellis

(Medical Xpress)—University of Adelaide researchers have found that women on antidepressant medication are more successful at breastfeeding their babies if they keep taking the medication, compared with women who quit antidepressants because of concerns about their babies' health.

These results have been presented this week at the 18th Perinatal Society of Australia and New Zealand (PSANZ) Annual Conference in Perth.

Using data from the Danish National Birth Cohort in Denmark, researchers in the University of Adelaide's Robinson Research Institute studied the outcomes of 368 women who were on [antidepressants](#) prior to becoming pregnant.

"We found that two thirds of the women (67%) stopped taking their antidepressant medication either after becoming pregnant or during breastfeeding," says NHMRC Early Career Fellow Dr Luke Grzeskowiak from the Robinson Research Institute.

"A third of the women (33%) continued to take antidepressant medication throughout their pregnancy and while breastfeeding, and these women were much more successful at maintaining breastfeeding up to and beyond the recommended six months.

"In contrast, those women who had stopped taking antidepressants were also more likely to stop breastfeeding within the recommended six

months."

Dr Grzeskowiak says the health benefits of continued breastfeeding greatly outweigh any perceived risk to the baby from antidepressant medication.

"This is a really important message because we know that breastfeeding has immense benefits for the child and the mum herself, including a degree of protection against post-natal depression," he says.

"The amount of antidepressant medication that finds its way into a mother's breast milk is very low. On the balance of it, we believe that continuing to take [antidepressant medication](#) and maintaining regular breastfeeding will be the best outcome for both the baby and the mother."

Dr Grzeskowiak says many [women](#) struggle with decisions about what to do with medications both during pregnancy and lactation.

"If they're taking antidepressants, they should be supported and encouraged by family members, friends and healthcare professionals to continue with their medication, knowing that good [breastfeeding](#) outcomes are all-important for them and their child," he says.

Provided by University of Adelaide

Citation: Proof that antidepressants and breastfeeding can mix (2014, April 10) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2014-04-proof-antidepressants-breastfeeding.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.