

Smoking visibility mapped for the first time

April 6 2014



The visibility of smoking in city streets has for the first time anywhere been mapped, in new research from the University of Otago, Wellington, New Zealand. Data from observations across the downtown area were mapped by the researchers, producing a record of the street areas where the most smokers could be seen. They used mapping methods previously used for landscape ecology and archeology. Credit: University of Otago, Wellington

The visibility of smoking in city streets has for the first time anywhere been mapped, in new research from the University of Otago, Wellington, New Zealand.

The research found that up to 116 smokers outside bars/cafés could be seen from any one location in the outdoor public areas of downtown Wellington (e.g. on a footpath). Of 2600 people observed in the outdoor areas of bars and cafés, 16% were [smoking](#), with a higher proportion than this in evenings.

Data from observations across the downtown area were mapped by the researchers, producing a record of the street areas where the most smokers could be seen. They used mapping methods previously used for landscape ecology and archeology.

Lead researcher Dr Amber Pearson says that the methods developed through this research will help policymakers demonstrate the visibility of smoking in different areas, and provide scientific evidence for [local authorities](#) to advance smokefree outdoor policies.

Another of the researchers, Associate Professor George Thomson, says the results show the need for policies to reduce the normality of smoking:

"Smokefree outdoor areas help [smokers](#) to quit, help those who have quit to stick with it, and reduce the normalisation of smoking for children and youth. They also reduce litter, water pollution and cleaning costs for local authorities and ratepayers," Thomson says.

In Australia, North America and other places, local authorities are increasingly creating smokefree streets and promoting smokefree al fresco dining and drinking, he says.

The study has been published in the international journal *BMC Public Health*.

Provided by University of Otago

Citation: Smoking visibility mapped for the first time (2014, April 6) retrieved 5 May 2024 from <https://medicalxpress.com/news/2014-04-visibility.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.