

20 years of data shows treatment technique improvement for advanced abdominal cancer

April 17 2014

Meaningful long-term survival is possible for selected patients suffering from advanced cancer of the abdomen when treated with cytoreductive surgery with Hyperthermic IntraPeritoneal Chemotherapy, or HIPEC, according to a first-of-its-size analysis by physicians at Wake Forest Baptist Medical Center.

Wake Forest Baptist has the largest reported, single-center experience with cytoreductive [surgery](#) and HIPEC, said lead author Edward A. Levine, M.D., and analysis of 20 years' worth of patient data shows that outcomes have clearly improved for [patients](#) undergoing this treatment technique.

Cytoreductive surgery, or debulking, is removal of part of a malignant tumor which can't be completely excised and is done to enhance [chemotherapy](#) effectiveness. HIPEC is a perfusion technique in which heated chemotherapy is administered directly into the abdomen during the surgery to kill remaining cancer cells.

The findings are published in the current issue of the *Journal of the American College of Surgeons* and were presented by Levine, professor of surgery and chief of surgical oncology service at Wake Forest Baptist, last year at the annual meeting of the Southern Surgical Association.

"Peritoneal dissemination of abdominal malignancy has a clinical course marked by bowel obstruction and death," Levine said. "It traditionally does not respond well to intravenous chemotherapy. This two-phase

approach has clearly extended the lives of many of our patients."

For the study, Levine and colleagues reviewed data of 1,000 patients who underwent 1,095 HIPEC procedures over the past two decades. The average age of patients was almost 53 years and slightly more than half were female. Primary tumor sites were: appendix 472(47.2 percent), colorectal 248(24.8 percent), mesothelioma 72(7.2 percent), ovary 69(6.9 percent), gastric 46(4.6 percent), others 97(9.7 percent).

Levine said the median overall survival was 29.9 months, with a five-year survival of 32.5 percent. "The data further shows that patients with advanced abdominal cancer should no longer be approached with therapeutic nihilism," he said. "Long-term survival is clearly possible for selected patients treated with HIPEC."

Provided by Wake Forest University Baptist Medical Center

Citation: 20 years of data shows treatment technique improvement for advanced abdominal cancer (2014, April 17) retrieved 13 May 2024 from <https://medicalxpress.com/news/2014-04-years-treatment-technique-advanced-abdominal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--