

## More access to health care may lead to unnecessary mammograms

May 27 2014

---



Credit: National Cancer Institute/public domain

Researchers have concluded that providing better access to health care may lead to the overuse of mammograms for women who regularly see a primary care physician and who have a limited life expectancy.

The cautionary note from researchers at the University of Texas Medical Branch at Galveston is that [screening](#) women in this category could

subject them "to greater risks of physical, emotional and economic suffering."

Dr. Alai Tan, a senior biostatistician in UTMB's Sealy Center on Aging and lead author of the study, said that "there has been little systematic attempt to define guidelines that would help determine when breast cancer screening might not be appropriate or overused.

"The American Cancer Society guidelines on screening, for example, have had no upper age limit," Tan wrote in the study. "This is different from the case with prostate-specific antigen screening, where both the American Cancer Society and the American Urological Association have longstanding guidelines that exclude men with a less than 10-year [life expectancy](#)."

The study was published in the June edition of *Medical Care*, the official journal of the Medical Care Section of the American Public Health Association.

Using data from 2006 through 2009, researchers studied about 5 percent the Medicare claims filed during that period by women whose life expectancy was less than seven years. They further studied where the women lived and whether they had a primary care physician.

In general, the researchers found that the use of mammograms decreases as a woman's life expectancy grows smaller. However, they found that the general downtrend as a woman ages could be offset by better access to [health care](#).

For example, the screening rate for woman with a life expectancy of less than six years who had seen a primary care physician two or fewer times during the three-year period studied was about 20 percent. However, among women who had seen a [primary care](#) physician 13 or more times

during that period who had a life expectancy of less than four years, the rate was about 34 percent.

The researchers also noted that there is about a four-year difference from when a cancer is diagnosed via screening and when it would be diagnosed clinically for women between 65 to 74. Previous studies have demonstrated that "survival benefits" were not apparent seven to 10 years after a screening.

The study also found that [women](#) with a limited life expectancy who lived in an area with greater access to mammography resources, more [primary care physicians](#), mammographic facilities and radiologists, were more likely to be screened.

Mammograms have been heavily promoted for about 25 years and many patients have come to expect that a screening test such as this would be a routine procedure, the researchers wrote. The researchers concluded that life expectancy should be a factor in deciding whether a mammogram is needed.

Provided by University of Texas Medical Branch at Galveston

Citation: More access to health care may lead to unnecessary mammograms (2014, May 27) retrieved 17 July 2024 from <https://medicalxpress.com/news/2014-05-access-health-unnecessary-mammograms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--