

Air pollution exposure in second trimester may increase asthma risk in children

May 19 2014

Children who are exposed in utero to high levels of particulate air pollution during the second trimester of pregnancy may be at greater risk of developing asthma in early childhood, according to a new study presented at the 2014 American Thoracic Society International Conference.

"We know that mothers' exposure to [air pollution](#) during pregnancy can affect lung development of their babies and lead to subsequent respiratory disorders, including asthma, although little is known about whether timing of the exposure is important to consider," said lead author Yueh-Hsiu Mathilda Chiu, ScD, from the Department of Pediatrics at Icahn School of Medicine at Mount Sinai in New York. "In our study, we assessed whether higher exposure to particulate air pollution at more specific time windows in pregnancy were particularly linked to higher asthma risk in urban children."

The study included 430 full-term children followed to age 7 years and their mothers. Daily exposure to air pollution from sources including traffic, power plants, and other industrial sources consisting of [fine particles](#) in the prenatal period was estimated based on where these mothers lived. These fine particles, which are more likely to be inhaled deep into the lungs, have been linked to the greatest health risk and previous studies have suggested that effects on pregnant women can be transferred to the growing baby.

The researchers found that exposure to higher levels of fine particles in

the second trimester was most strongly associated with increased asthma onset among the children, particularly for those born to non-obese mothers. "It is possible that the effect of maternal obesity, another known risk factor of childhood asthma onset, may be so strong that it was difficult to determine additional effects of air pollution among children born to obese mothers in this setting."

"While we should continue to improve air quality and minimize exposure to [pregnant women](#) throughout the entire pregnancy for a host of health reasons," said Dr. Rosalind Wright, MD MPH, senior investigator from the Department of Pediatrics at Icahn School of Medicine at Mount Sinai, "pinpointing the gestational period during which air pollution has the greatest effects on the developing lung may add to our understanding of the mechanisms underlying this relationship."

More information: Abstract 53288, Identifying Prenatal Windows Of Susceptibility To Particulate Air Pollution On Childhood Asthma Onset In A Prospective Urban Birth Cohort , Scientific Abstract , 06.01 - Air Pollution: Epidemiology and Mechanisms (EOH) , Y.-H.M. Chiu¹, B.A. Coull², I. Kloog², J. Schwartz², H.-H.L. Hsu¹, R.O. Wright¹, R.J. Wright¹; ¹Icahn School of Medicine at Mount Sinai - New York, NY/US, ²Harvard School of Public Health - Boston, MA/US

Provided by American Thoracic Society

Citation: Air pollution exposure in second trimester may increase asthma risk in children (2014, May 19) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-05-air-pollution-exposure-trimester-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.